

List of Tables

1	Chemical Fingerprinting of Spilled or Discharged Petroleum — Methods and Factors Affecting Petroleum Fingerprints in the Environment	
1-1	Average Estimated Annual Releases (1990–1999) of Petroleum into the Marine Environment in North America and worldwide by Source (in Thousands of Metric Tonnes). Data from National Research Council (2002)	2
1-2	PAH and Related Compounds or Groups Commonly Targeted in Oil Spill-Related Samples Obtained Using GC/MS-SIM	8
1-3	Inventory of Petroleum Biomarkers Commonly Used in the Characterization of Spilled Oil Obtained Using GC/MS-SIM	9
1-4	Inventory of the Factors Controlling the Chemical Fingerprints of Spilled or Discharged Petroleum	11
1-5	Inventory of Common Biomarker Compound Classes Available for Chemical Fingerprinting of Petroleum in the Environment. The Table Includes the Carbon Boiling Range over Which Each Class Occurs and the Dominant Mass/Charge (m/z) Ratio upon Which They Can Be Identified Using Gas Chromatography-Mass Spectrometry	16
1-6	Typical Distillation Fractions and Petroleum Products Produced from Crude Oil. (Boiling and Carbon Ranges Are Approximate and Can Vary Slightly with Refiner)	22
2	Spill Site Investigation in Environmental Forensic Investigation	
2-1	Characteristics of Spills on Land and Water (Adapted from Owens, 2002)	56
2-2	Examples of Factors to be Assessed as Part of a Site Investigation	57
2-3	Priorities of the Safety Officer during an Oil Spill (Adapted from IPIECA, 2002)	58
3	Petroleum Biomarker Fingerprinting for Oil Spill Characterization and Source Identification	
3-1	Petroleum Biomarkers Frequently used for Forensic Oil Spill Studies	78
3-2	Common Modifiers and Nomenclatures Used to Modify the Structural Specification of Cyclic Biomarkers	79
3-3	Surrogate and Internal Standard Used for Oil and Biomarker Fingerprinting	83
3-4	Characteristic Fragment Ions for Various Biomarkers	92
3-5	Quantitation results of Major Target Biomarkers in Example Crude Oils and Petroleum Products	98
3-6	Quantitation Results of Target Biomarkers in the MESA Oil and Its Four Fractions	107
3-7	Peak Identification of the Triaromatic and Monoaromatic Steranes in the Platform Elly Oil	109
3-8	Peak Identification of Diamondoid Compounds in Prudhoe Bay Oil	116

3-9	Diagnostic Biomarker Ratios Frequently used for Identification, Correlation, and Differentiation of Spilled Oils	127
3-10	Quantitation Results and Diagnostic Ratios of Sesquiterpanes in Three Oil Samples	137
4	Characterization of Polycyclic Aromatic Sulfur Heterocycles for Source Identification	
4-1	Names of Sulfur Compounds in Petroleum as Illustrated in Figure 4-1	149
4-2	Polycyclic Aromatic Sulfur Heterocycles Identified in the Egyptian Crude Oil in Figure 4-2	152
5	Oil Spill Identification by Comprehensive Two-Dimensional Gas Chromatography (GC × GC)	
5-1	Peak identities of Compounds Labeled in Figure 5-24	201
6	Application of Stable Isotope Ratios in Spilled Oil Identification	
6-1	Stable Isotope and Metal Concentrations in Spilled Oils Near a Tank Farm in Western U.S. Compared with Potential Crude Oil Sources	213
6-2	Sensitivity and Precision for Elements Reported to Date of Compound-Specific Isotope Analysis (CSIA). Data from Schmidt et al. (2004)	214
7	Emerging CEN Methodology for Oil Spill Identification	
7-1	Diagnostic Ratios Derived from Selected <i>n</i> -Alkanes and Acyclic Isoprenoids (Diterpanes) Recorded by GC/FID	236
7-2	Recommended Diagnostic Ratios (DR) Derived from Alkylated PACs	237
7-3	Recommended Diagnostic Ratios (DR) Derived from Tri- and Pentacyclic Triterpanes	242
7-4	Recommended Diagnostic Ratios (DR) from Rearranged (Diasteranes) and Regular 14 α (H)- and 14 β (H)-Steranes	243
7-5	Recommended Diagnostic Ratios (DR) Derived from Triaromatic Steroids	243
7-6	Optional Diagnostic Ratios (DR) Derived from Sesquiterpanes	244
7-7	Example of Comparison of Diagnostic Ratios Based on a Repeatability Limit of 14%	248
7-8	Comparison of Diagnostic Ratios of Acyclic Isoprenoids Using the Repeatability Limit	253
7-9	Comparison of Diagnostic Ratios (in %) of Some Triterpanes Using the Repeatability Limit	253
7-10	Comparison of Diagnostic Ratios (in %) of Spill and Candidate Source Samples Using a Repeatability Limit ($r_{95\%}$) of 14%	254
8	Advantages of Quantitative Chemical Fingerprinting in Oil Spill Identification	
8-1	Inventory of Target Saturated Hydrocarbons (SHC), PAHs, and Biomarkers Commonly Analyzed by GC/FID and GC/MS-SIM in Oil Spill Studies	267
8-2	Summary of Typical Data Quality Objectives Used in Quantitative Fingerprinting Studies of Waterborne Oil Spills	269
8-3	List of Common Quantitative Indices Useful in Oil Spill Investigations. This List Is Not Intended to Be All Inclusive	271
8-4	Diagnostic Ratios and Precision-Based Statistics for a Spilled Oil and Two Candidate Source Oils (A and B). All Data are Based Upon the Absolute	

	Concentrations of Individual Compound or Compound Groups Measured by GC/MS as per the Revised Nordtest Method (Daling et al., 2002)	272
8-5	Example Data for Selected Biomarkers for Two Oils Demonstrating How the Absolute Analyte Concentrations are Needed to Calculate Accurate Diagnostic Ratios in Theoretical Mixtures. Ratios Calculated from the Original Ratios, as Might Be Available from Peak Areas (but not Concentrations), Do Not Mix Linearly Due to Concentration Differences in the Two Oils. Therefore, Absolute Concentrations Are Necessary to Develop Accurate Mixing Models	277
8-6	Diagnostic Ratios for the IFO and HFO Source Oils and Field Samples	282
9	A Multivariate Approach to Oil Hydrocarbon Fingerprinting and Spill Source Identification	
9-1	List of Mass Fragments (m/z) and Corresponding Compound Groups Analyzed with High Relevance for Oil Hydrocarbon Fingerprinting	300
9-2	Application of a Variable-Outlier Detection Method to 17 Biomarker Ratios	313
10	Chemical Heterogeneity of Marine Heavy Fuel Oils	
10-1	ISO Specifications for Marine Residual Fuel Oils	329
10-2	Weight Percentages of Total Hydrocarbons (THC) and Common Isoalkane Ratios for 71 Worldwide IFO 380 Heavy Fuel Oils	331
10-3	Homo- and Heteroatomic PAH Composition for 71 Worldwide IFO 380 Heavy Fuel Oils (Concentrations in $\mu\text{g}/\text{kg}$)	339
10-4	Summary Statistics for Certain PAH Diagnostic Metrics for 71 Worldwide IFO 380 Heavy Fuel Oils (See Table 10-3 for Compound Abbreviations.)	342
10-5	Biomarker-Based Thermal Maturity Parameters Measured in Five Asphalts (S_1 – S_5) Manufactured with Vacuum Distillation Residuum Derived from the Parent Crude Oil Source	345
11	Biodegradation of Oil Hydrocarbons and Its Implications for Source Identification	
11-1	Comparison of Aerobic and Anaerobic Respiration Reactions	366
13	Trajectory Modeling of Marine Oil Spills	
13-1	Grid Resolution of Atmospheric Models Modified from Kalnay (2003)	411
13-2	Estimated Length and Time Scale of Oil Movement Due to Surface Current Transport	411
14	Oil Spill Remote Sensing: A Forensics Approach	
14-1	Relationships between Appearance and Oil-Slick Thickness	430
14-2	Relationship between Oil Thickness and Detection Limits in Remote Sensing Instruments	432
14-3	Attributes for Sensor Selection	438
14-4	Sensor Suitability for Various Missions	439
15	Advances in Forensic Techniques for Petroleum Hydrocarbons: The <i>Exxon Valdez</i> Experience	
15-1A	Major Hydrocarbon Fractions, TOC, and PAH Characteristics of EVC, Weathered EVC, PWS Prespill Benthic Sediments, Oil Residues at Historical Human Activity (HA) Sites, and Potential Background Sources	455

15-1B	Saturate and Aromatic Biomarker Indices for EVC, Weathered EVC, PWS Prespill Benthic Sediments, Oil Residues at Historical Human Activity (HA) Sites, and Potential Background Sources	456
15-2	Summary of Source Allocation Results for Subtidal Sediments Sampled in Six PWS Embayments in 1999 and 2000	471
15-3	Interpretations of Tissue (Excluding Shellfish) PAH Compositions Reported in the 2004 Version of EVTHD (See Text for Definition of EVTHD)	475
16	Case Study: Oil Spills in the Strait of Malacca, Malaysia	
16-1	PAHs and alkanes compositions for crude oils and Malaysian tar-ball	500
17	Evaluation of Hydrocarbon Sources in Guanabara Bay, Brazil	
17-1	Results for the Individual PAH (ng/g dry weight)* of Sediment Samples from Guanabara Bay — Campaign 2000 and Studied Oils	512
17-2	Results for the Individual PAH (ng/g dry weight)* of Sediment Samples from Guanabara Bay — Campaign 2003 and Studied Oils	514
17-3	Literature Data on PAH Concentration (ng/g dry weight) of Sediments from Various Coastal Sites in the World	516
17-4	Results for Biomarkers, Terpanes, and Steranes of Sediment Samples from Guanabara Bay — Campaign 2000	528
17-5	Results for Biomarkers, Terpanes, and Steranes of Sediment Samples from Guanabara Bay — Campaign 2003 the 30-Norhopane Series	530