

List of Figures

1	Chemical Fingerprinting of Spilled or Discharged Petroleum — Methods and Factors Affecting Petroleum Fingerprints in the Environment	
1-1	Diagram depicting the four factors that affect the chemical fingerprints of petroleum relevant in oil spill investigations.	4
1-2	Ternary diagram showing the general compositional features of crude oils. After Tissot and Welte (1984). Arrow indicates general compositional change due to biodegradation.	13
1-3	Examples of aliphatic and aromatic hydrocarbons in crude oils.	14
1-4	Example of the disparate primary chemical fingerprinting features in crude oils from the North Slope of Alaska and the Niger Delta, Nigeria. (a–b) GC/FID chromatograms, (c–d) partial m/z 191 mass chromatograms, and (e–f) partial m/z 217 mass chromatograms. * = internal standards, left to right: <i>o</i> -terphenyl, 5α -androstane and tetracosane- d_{50} .	17
1-5	GC/FID chromatograms and partial m/z 191 mass chromatograms for a mystery oil spill and produced crude oil from two nearby production platforms. Differences in the relative abundance of gammacerane were sufficient to distinguish these (and other) genetically related oils from one another. * = internal standards.	19
1-6	Summary of foreign and domestic petroleum products transported as cargo in U.S. waterways and ports in 2003. Volume calculated from short ton using average density. Data from WCUS (2003).	19
1-7	Schematic flowchart of a modern complex refinery highlighting the distillation, conversion, and finishing processes used in the production of petroleum products.	21
1-8	Gas chromatography-flame ionization detection (GC/FID) chromatograms of distinct petroleum products. (A) automotive gasoline, (B) jet fuel A, (C) diesel fuel #2, (D) heavy fuel oil #6, (E) lubricating oil, and (F) petroleum asphalt. # = <i>n</i> -alkane carbon number.	23
1-9	Partial m/z 191 mass chromatograms for a (A) parent crude oil feedstock and (B) daughter distillate cut showing the progressive loss of tricyclic triterpanes due to distillation effects.	25
1-10	Partial m/z 123 mass chromatograms showing the distributions of sesquiterpanes in (A) a spilled jet fuel and (B) a candidate source oil. Peak numbers correspond to compounds described by Alexander et al. (1984). The large peak between peaks 10 and 1 is unidentified.	25
1-11	Chemical fingerprints for two distillate fuel blending stocks. (A) GC/MS TIC for straight-run (virgin) gas oil, (B) GC/MS TIC for light cracked distillate (cycle oil), (C) PAH histogram for straight-run gas oil, and (D) PAH histogram for light cracked distillate. # = <i>n</i> -alkane carbon number, Pr = pristane, Ph = phytane; for PAH analytes, see Table 1-2.	26

1-12	GC/FID chromatograms for four base oils used in the manufacture of lubricating oils. (A) low-viscosity base oil (70N), (B) medium-viscosity base oil (150N), (C) heavy-viscosity base oil (320N), and (D) bright stock (850N).	29
1-13	GC/FID of a bilge oil containing a mixture of different petroleum. (Modified from Emsbo-Mattingly, 2006.)	30
1-14	GC/FID chromatograms showing the progressive loss of compounds from a 34°API crude oil spilled following Hurricane Katrina. (A) slightly evaporated crude oil, (B) moderately evaporated crude oil, and (C) severely evaporated crude oil.	31
1-15	GC/FID chromatograms showing the effect of wax precipitation on the chemical fingerprint of a paraffinic oil. (A) starting oil, (B) wax-depleted fraction within spilled oil 71 hours after the spill, and (C) wax-enriched fraction within the spilled oil 71 hours after the controlled spill experiment. Data from Daling et al. (2002).	35
1-16	GC/FID chromatogram of the TPH extracted from a Louisiana peat containing 2000 mg/kg (dry).	39
1-17	Histograms showing the distribution of PAH in three U.S. coals of varying rank and in modern wood-derived charcoal. For PAH compound abbreviations, see Table 1-2.	40
1-18	GC/FID fingerprints showing the chromatographic character of the UCM “humps” in (A) urban sediment impacted by urban runoff, (B) motor oil, and (C) hydraulic oil. OTP (<i>o</i> -terphenyl) and 5 α -androstane are internal standards.	42
1-19	Histograms showing the distributions and concentrations of PAH in an urban sediment from the Elizabeth River (Virginia) containing anthropogenic background PAH and in various urban runoff candidate source components. For compound abbreviations, see Table 1-2. Concentrations refer to mg of PAH per kg of sediment (dry), urban dust (dry), soot (dry), oil, and asphalt (dry).	44
2	Spill Site Investigation in Environmental Forensic Investigation	
2-1	Map of tar ball distribution indicating those that matched the source oils on the vessel and those that did not match (adapted from Owens et al., 2002).	69
2-2	Example of chain-of-custody form.	70
3	Petroleum Biomarker Fingerprinting for Oil Spill Characterization and Source Identification	
3-1	Molecular structures of example cyclic terpenoid compounds in oil.	76
3-2	Types of isomers.	80
3-3	Oil sample preparation flowchart.	82
3-4	Mass spectra of some common biomarkers used in environmental forensic studies.	87
3-5	The SIM chromatograms at <i>m/z</i> (a) 191 and (b) 217 of a Kuwait crude oil for common terpane and sterane biomarker classes.	91
3-6	Carbon range of common cyclic biomarker classes in crude oil and petroleum products.	93
3-7	GC-FID (left panel) and GC-MS (at <i>m/z</i> 191, right panel) chromatograms of five different oils (light to medium) to illustrate differences in the <i>n</i> -alkane and tri-, tetra-, and pentacyclic terpane distributions between oils.	94

3-8	GC-MS chromatograms at m/z 217 and 218 for five different oils (light to medium) to illustrate differences in sterane distributions between oils.	95
3-9	GC-FID (left panel) and GC-MS (at m/z 191, right panel) chromatograms of five heavy oils from different regions.	96
3-10	GC-MS chromatograms at m/z 217 and 218 of five heavy oils from different regions.	97
3-11	GC-FID (left panel) and GC-MS (at m/z 191, right panel) chromatograms of five common petroleum products (light to heavy) to illustrate differences in the n -alkane and tri-, tetra-, and pentacyclic terpane distributions between oils.	100
3-12	GC-MS chromatograms at m/z 217 and 218 for common petroleum products (light to heavy) to illustrate differences in sterane distributions between oils.	102
3-13	GC-FID (left panel) and GC-MS (at m/z 191, middle panel; m/z 218, right panel) chromatograms of five lubricating oils.	103
3-14	GC-FID chromatograms (left panel) and PAH distributions (right panel) of the MESA oil and its four fractions.	105
3-15	GC-MS chromatograms of the MESA oil and its four fractions at m/z 191 (left panel) and 218 (right panel) to illustrate differences in biomarker distributions.	106
3-16	Peak identification of the triaromatic (m/z 231) and monoaromatic (MA, m/z 253) steranes in the Platform Elly oil.	108
3-17	Mass chromatograms of the TA- (m/z 231) and MA-steranes (253) in the aromatic hydrocarbon fractions of the NIST 1582 oil, a Diesel No. 2 from Korea, IFO-180, and a lubricating oil.	110
3-18	GC-MS (SIM) chromatograms of sesquiterpanes eluting in the n -C ₁₃ and n -C ₁₆ range.	112
3-19	<i>Left panel:</i> GC-MS chromatograms of sesquiterpanes at m/z 123 for light (API > 35), medium (API: 25–35), and heavy (API < 25) crude oils, including Alaska North Slope (ANS), Arabian Light, Scotia Light oil (Nova Scotia), West Texas, and California API 11. <i>Right panel:</i> Comparison of SIM chromatograms of sesquiterpanes at m/z 123 for common petroleum products.	113
3-20	GC-MS chromatograms of diamondoids in Prudhoe Bay oil for peak identification.	115
3-21	GC-MS total ion chromatograms of adamantanes (left) and diamantanes (right) of five representative crude oils including Cold Lake Bitumen, Maya, Mars, Alaska North Slope, and South Louisiana crude oil.	118
3-22	GC-MS total ion chromatograms of adamantanes and diamantanes of five representative refined petroleum products including Jet A, Diesel No. 2, Korea Diesel, Fuel No. 4, and Valvoline-10W-30 lube oil.	119
3-23	Comparison of GC-FID and GC-MS (m/z 191) of three unknown oil samples.	122
3-24	Molecular structures of a selection of “source-specific” biomarkers.	123
3-25	A: Cross-plots of the double ratios of Peak 4/Peak 5 versus Peak 3/Peak 5 for over 50 different oils and refined products. The circle indicates related samples from the same origin. B: Cross-plots of the double ratios of Peak 4/Peak 5 versus Peak 3/Peak 5 for 11 weathering oil series and 1 diesel weathering series. Each weathering oil series produces a tight cluster.	129
3-26	Extracted ion chromatograms at m/z 123, sesquiterpanes, for three Round Robin samples (right) and their corresponding GC-FID chromatograms for n -alkane analysis (left).	136

- 3-27 Correlation of diagnostic ratios (normalized to %) of sesquiterpanes between spill sample 2 and suspected source samples 1 (left) and 3 (right) at 95% confidence. All the data (A: left panel) overlap the 1:1 line at 95% confidence, representing a “perfect” match between samples 2 and 1. Conversely, most data points (B: right panel) between samples 2 and 3 do not overlapping the line at 95% confidence, representing a “nonmatch.” 138
- 3-28 Comparison of GC-MS chromatograms of terpanes (m/z 191) and steranes (m/z 218) in three oil samples. 139
- 4 Characterization of Polycyclic Aromatic Sulfur Heterocycles for Source Identification
- 4-1 Structures of sulfur compounds in petroleum. For names, see Table 4-1. 149
- 4-2 Partial extended AED chromatogram of polyaromatic fraction showing the distribution and relative retention times of PASH in an Egyptian crude oil. Peak identifications are listed in Table 4-2. 5FBT, 2FDBT, and 6FBN[2,3-d]T are fluorinated internal standards. (Reprinted from Hegazi et al., 2003, with permission from Elsevier Science.) 151
- 4-3 GC-FID chromatograms of (a) nondesulfurized diesel fuel and (b) deeply hydrodesulfurized diesel fuel. The asterisk indicates an impurity from the dichloromethane used in the sample workup. (Reprinted from Schade et al., 2002, with permission from Taylor & Francis.) 153
- 4-4 Kendrick mass defects versus Kendrick nominal mass of vacuum residues: (a) first ligand exchange chromatography (LEC) fraction, (b) second LEC fraction before HDS, (c) first LEC fraction, and (d) second LEC fraction after HDS. The double bond equivalent (DBE) scale given is valid for both panels. (Reprinted from Müller et al., 2005, with permission from the American Chemical Society.) 154
- 4-5 Gas chromatogram of a diesel (350 ppm sulfur) monitored at m/z 184, which is the mass for DBT. Coinjection with DBT reveals this compound at a retention time of 16.86 minutes. 158
- 4-6 Mass spectrum of the peak at 16.86 minutes in Figure 4-5 (top) and a mass spectrum of pure DBT (bottom). 159
- 4-7 Plot of the relative ratios of 2-/3-methyl-DBT to 4-methyl-DBT versus the relative ratios of 1-methyl-DBT to 4-methyl-DBT for 25 different oils. The circles 1 and 2 indicate related samples from origins of North Slope and Terra Nova, respectively. (Reprinted from Wang and Fingas, 1995, with permission from the American Chemical Society.) 160
- 4-8 Mass fragmentograms of methyl-DBTs (m/z 198) from clams collected in April 1993 (lower trace) and December 1995 (upper trace). 2- and 3-MDBT are biodegraded appreciably faster than 1- and 4-MDBT. (Reprinted from Albaigés et al., 2000, with permission from WIT Press.) 163
- 4-9 Double-ratio plot of C2/C3 dibenzothiophenes versus C2/C3 phenanthrenes from oil residues from Prince William Sound, Alaska, eight years after the *Exxon Valdez* spill. NSC is the source oil from the *Exxon Valdez*. (Reprinted from Michel and Hayes, 1999, with permission from Elsevier Science.) 163
- 5 Oil Spill Identification by Comprehensive Two-Dimensional Gas Chromatography
- 5-1 Conceptual diagram of matching mixture dimensions with separation dimensions to achieve complete separation and an ordered chromatogram.

- (A) Three-dimensional mixture in gray scale, shape, and size; (B) one-dimensional separation in size results in poor separation; (C) two-dimensional separation improves separation and results in a more ordered chromatogram; (D) matching mixture and separation dimensions results in complete separation and ordered chromatogram. 170
- 5-2 Concepts of comprehensive two-dimensional gas chromatography ($GC \times GC$): (A) first-dimension chromatogram peak containing coelutents; (B) modulator samples first-dimension peaks and injects narrow peak into second dimension; (C) coelutents are separated in the second dimension; (D) serial flame ionization detector data stream is sliced into segments and stacked into an array; (E) array is visualized with a contour. Copyright 2002 from *GC \times GC — A New Analytical Tool for Environmental Forensics*, by G.S. Frysinger et al. Reproduced by permission of Taylor & Francis Group, LLC, <http://www.taylorandfrancis.com>. 171
- 5-3 Chromatograms of *Exxon Valdez* cargo oil: (A) One-dimensional gas chromatogram using a nonpolar 100% polydimethylsiloxane stationary phase; (B) $GC \times GC$ volatility-by-polarity interpolated color contour plot. The first dimension is a separation using a nonpolar 100% polydimethylsiloxane stationary phase and the second dimension is a separation using a polar 50% phenyl polysilphenylene stationary phase; (C) a small portion of the $GC \times GC$ chromatogram visualized as a mountain plot. The mountain plot is excellent for visualizing relative differences among neighboring peaks. See color plate. 174
- 5-4 A small portion of a volatility-by-polarity $GC \times GC$ chromatogram of *Exxon Valdez* cargo oil showing the region containing the triaromatic sterane, hopanes, and sterane biomarkers. GC-MS extracted ion chromatograms (EICs) of these biomarker target ions are overlaid for identification and comparison. Peaks numbered on each EIC are identified in Frysinger and Gaines (2001). The structures of $5\alpha,14\alpha,17\alpha(H)$ -cholestane (20R) (peak 46), $17\alpha,21\beta(H)$ -hopane (peak 58) and cholestane (20S) (peak 69) are included for reference. *x*-axis: minutes; *y*-axis: seconds. Reproduced with permission from Frysinger and Gaines (2001). Copyright 2001 Wiley-VCH. 176
- 5-5 Partial $GC \times GC$ -MS chromatogram of diesel fuel. The *x*-axis is a volatility-based chromatographic separation on a 100% polydimethylsiloxane stationary phase. The *n*-alkanes decane (C_{10}) to tridecane (C_{13}) are identified. The *y*-axis produces polarity-based chromatographic separation on a 14% cyanopropylphenyl methylsiloxane stationary phase. The MS total ion intensity data are displayed as an interpolated color contour. Overlay data: (A–B) GC-MS spectra for specific retention times; (C) GC-MS *m/z* 120 extracted ion chromatogram; (D–H) $GC \times GC$ -MS spectra and structures of naphthalene, 1,2,3-trimethylbenzene, 4-methyldecane, *n*-butylcyclohexane, and decahydronaphthalene, respectively; (I) indan; (J) indene. Copyright 2002 from *GC \times GC — A New Analytical Tool for Environmental Forensics*, by G.S. Frysinger et al. Reproduced by permission of Taylor & Francis Group, LLC, <http://www.taylorandfrancis.com>. 177
- 5-6 Effect of changing the second-dimension stationary phase on the separation of nonpolar saturates: (A) $GC \times GC$ chromatogram of the saturates fraction of a sediment extract using a polar second-dimension 14% cyanopropylphenyl polysiloxane stationary phase; (B) $GC \times GC$ chromatogram of the same saturates fraction using a chiral γ -cyclodextrin second-dimension

- stationary phase. The expected positions of the n -C₁₃ to n -C₂₀ alkanes are marked with circles. The region inside the dotted lines is expanded in (C); (C) expanded region showing expected position of n -C₁₅ and location of branched alkane bands. Reprinted with permission from Frysinger et al., 2003. Copyright 2003, American Chemical Society. See color plate. 179
- 5-7 Location of petroleum chemical classes on a volatility-by-polarity GC × GC chromatogram. Nonpolar alkanes are located at the bottom, with one- and multiring aromatics further up the y -axis as their polarity increases. Inset A: naphthalene region showing bands of alkyl-substituted naphthalenes and biphenyls; Inset B: nonpolar region showing high-resolution separations of higher boiling (C₂₈₊) acyclic isoprenoids, normal and branched alkanes, and cyclic alkanes. See color plate. 180
- 5-8 Portion of a volatility-by-polarity GC × GC-FID chromatogram from each of the three samples compared: (a) spill; (b) and (c) potential spill sources. Both axes are in seconds. The boxes contain chemical families and individual compounds used to qualitatively and quantitatively compare two potential source samples with the spill samples. The first column separation shown along the x -axis was accomplished with a nonpolar 5% phenylmethylsiloxane stationary phase. The second column separation shown along the y -axis was accomplished with a polar polyethylene glycol/siloxane copolymer stationary phase. Reprinted with permission from Gaines et al., 1999. Copyright 1999, American Chemical Society. See color plate. 182
- 5-9 Quantitative comparisons between spill sample and two potential source samples from Figure 5-8. A quality-control (QC) sample, split from the spill, was also quantified. Each panel represents the results of integrating peaks within an individual box shown in Figure 5-8. Each bar represents the total peak or band of peaks integration results normalized to the first peak in the box. The error bar represents an estimated 10% variation calculated from replicate data, but is shown only in the plus direction. Compound classes represented: (a) alkylnaphthalene bands normalized to 2,3,5-trimethylnaphthalene; (b) alkylphenanthrenes normalized to phenanthrene; (c) cycloalkanes normalized to pentadecane; (d) heptadecane, octadecane, pristane, and phytane normalized to each other. Reprinted with permission from Gaines et al., 1999. Copyright 1999, American Chemical Society. 183
- 5-10 Map of the general area near the 1969 grounding of the barge *Florida*. 185
- 5-11 Gas chromatograms of Wild Harbor marsh sediment extracts: (A) Nov. 1973; (B) Aug. 2000. Reprinted with permission from Reddy et al., 2002. Copyright 2002, American Chemical Society. 185
- 5-12 Chromatograms of a Wild Harbor sediment extract, Aug. 2000: (A) Conventional one-dimensional gas chromatogram. The separation was accomplished using a nonpolar polydimethylsiloxane stationary phase; (B) volatility-by-polarity GC × GC chromatogram. The first column separation along the x -axis was accomplished using a nonpolar polydimethylsiloxane stationary phase. The second column separation was accomplished using a polar 14%-cyanopropylphenyl polysiloxane stationary phase. Reprinted with permission from Reddy et al., 2002. Copyright 2002, American Chemical Society. See color plate. 186
- 5-13 Comparison of Wild Harbor sediment extract with weathered MERL oil: (A) Enlarged volatility-by-polarity GC × GC chromatogram of Wild Harbor extract, Aug. 2000 (as in Figure 5-12); (B) neat MERL oil; (C) 30% by mass

- weathered (WX) MERL oil; (D) 70% by mass weathered (WX) MERL oil. The first column separation along the x -axis was accomplished using a nonpolar polydimethylsiloxane stationary phase. The second column separation was accomplished using a polar 14%-cyanopropylphenyl polysiloxane stationary phase. Weathering experiments were performed in a hood where neat MERL oil was allowed to evaporate until the desired mass loss was achieved. The background is blue. Peak intensity is scaled from white, to red, and then to blue (most intense). Compound abbreviations: (N) naphthalene; (C1N) C₁-naphthalenes; (C2N) C₂-naphthalenes; (C3N) C₃-naphthalenes. Reprinted with permission from Reddy et al., 2002. Copyright 2002, American Chemical Society. See color plate. 188
- 5-14 Map of the general area near the grounding of the barge *Bouchard 65*. The grounding of the barge *Florida*, discussed previously, is also shown for reference. 189
- 5-15 Chromatograms of the sediment extract taken from Winsor Cove in 2002 approximately 30 years after being contaminated with No. 2 fuel oil: (A) one-dimensional gas chromatogram showing UCM and the region of the chromatogram visualized in (B). The separation was accomplished using a polydimethylsiloxane stationary phase; (B) volatility-by-polarity GC \times GC-FID chromatogram showing the region containing the target compounds. The first-column separation shown along the x -axis was accomplished with a polydimethylsiloxane stationary phase. The second-column separation shown along the y -axis was accomplished with a 50%-phenylpolysilphenylene-siloxane stationary phase. See color plate. 190
- 5-16 Possible structure of C₅- and C₆-decalins found in degraded No. 2 fuel oil in Winsor Cove approximately 30 years after a spill: (A) 8 β (H)-drimane; (B) 8 β (H)-homodrimane. 190
- 5-17 Gas chromatograms of *Bouchard 65* incubation sediment extracts artificially degraded in the laboratory: (A) day 1; (B) day 11; (C) day 46. 191
- 5-18 Volatility-by-polarity GC \times GC $m/z = 123$ extracted ion chromatographs of *Bouchard 65* incubation sediment extracts showing the C₄- to C₆-decalin region: (A) neat *Bouchard 65* fuel oil; (B) day 11; (C) day 36; (D) day 46; (E) day 46 spiked with cis- and trans-1,1,4,4,6-pentamethyldecalin standard; (F) Winsor Cove sediment extract showing the bands of peaks containing C₄-decalins (C4D), C₅-decalins (C5D), C₆-decalins (C6D), and location of 8 β (H)-drimane (A) and 8 β (H)-homodrimane (B) whose structures are given in Figure 5-16. The first-column separation shown along the x -axis was accomplished with a nonpolar polydimethylsiloxane stationary phase. The second-column separation shown along the y -axis was accomplished with a polar 50%-phenylpolysilphenylene-polysiloxane stationary phase. See color plate. 192
- 5-19 Map of Buzzards Bay showing the track of the *Bouchard 120* barge on April 27, 2003. Oil-impacted beaches are highlighted in gray around the perimeter of the bay. Nyes Neck (where the oil spill samples discussed in this section were collected) is shown on the inset map. Copyright 2006 from *Tracking the Weathering of an Oil Spill with Comprehensive Two-Dimensional Gas Chromatography*, by Nelson et al. Reproduced by permission of Taylor & Francis Group, LLC., <http://www.taylorandfrancis.com>. 193
- 5-20 Conventional GC and GC \times GC chromatograms of sample extracts from *Bouchard 120* No. 6 fuel oil-covered rocks collected at various times after

- the spill: (A) conventional GC chromatogram of sample taken 12 days after the spill. The separation was achieved using a polydimethylsiloxane stationary phase; (B) GC × GC volatility-by-polarity chromatogram of sample taken 12 days after the spill. The first dimension separation along the *x*-axis was achieved using a nonpolar polydimethylsiloxane stationary phase. The second dimension separation along the *y*-axis was achieved with a polar 50% phenylpolysilphenylenesiloxane stationary phase; (C) conventional GC chromatogram of sample taken 179 days after the spill; (D) GC × GC volatility-by-polarity chromatogram of sample taken 179 days after the spill; (E) a point-by-point difference chromatogram produced by the subtraction of chromatogram (D) from chromatogram (B) after normalization to the conserved standard 17 α (H)-21 β (H)-hopane (marked with a star). Copyright 2006 from *Tracking the Weathering of an Oil Spill with Comprehensive Two-Dimensional Gas Chromatography*, by Nelson et al. Reproduced by permission of Taylor & Francis Group, LLC., <http://www.taylorandfrancis.com>. 194
- 5-21 Percent losses of select hydrocarbons from *Bouchard 120* No. 6 fuel oil covered rocks collected at Nyes Neck after six months of weathering: (●) *n*-alkanes; (◆) *n*-alkylcyclohexanes; (△) branched alkanes. Copyright 2006 from *Tracking the Weathering of an Oil Spill with Comprehensive Two-Dimensional Gas Chromatography*, by Nelson et al. Reproduced by permission of Taylor & Francis Group, LLC., <http://www.taylorandfrancis.com>. 196
- 5-22 Volatility-by-polarity GC × GC chromatogram of (A) reservoir crude oil from platform Holly, Santa Barbara Channel, California, and (B) oil stringer emerging through the sea floor collected at the Jackpot seep field, Santa Barbara Channel, California. The first-column separation shown along the *x*-axis was accomplished with a nonpolar polydimethylsiloxane stationary phase. The second-column separation shown along the *y*-axis was accomplished with a polar 50%-phenylpolysilphenylene-siloxane stationary phase. A star on each chromatogram indicates the position of the internal standard dodecahydrotriphenylene (DDTP). The distinct difference in *n*-alkane content between the two samples is evident in this figure. See color plate. 197
- 5-23 Partial volatility-by-polarity GC × GC chromatogram of the platform Holly produced crude oil showing peaks in the vicinity of *n*-C₃₃ through *n*-C₃₉. The location of high-molecular-weight isoprenoid alkanes (both acyclic and cyclic), thought to be biomarkers of Archaeal membrane lipids, are indicated along with some representative molecular structures. See color plate. 199
- 5-24 Partial volatility-by-polarity GC × GC chromatogram of the platform Holly reservoir crude oil showing peaks in the vicinity of the sterane and desmethylsterane biomarkers. Identification of labeled peaks is found in Table 5-1. See color plate. 200
- 6 Application of Stable Isotope Ratios in Spilled Oil Identification
- 6-1 Carbon isotope ratios in a variety of geochemical materials. (Reprinted with permission from Fuex (1977). Copyright (1977) Elsevier Science.) 209
- 6-2 Schematic diagram of an instrument set-up to perform compound specific carbon isotope analysis. 209
- 6-3 Carbon isotope ratios of Exxon Valdez oil and residues, and other tar balls collected in Prince William Sound, Alaska and analyzed by Kvenvolden et al. (1997). 211

6-4	Carbon isotope ratios of saturate and aromatic fractions of oil and tar balls associated with the BP American Trader spill offshore Huntington Beach, California. These are compared with ratios of tar balls north and south of the spill site.	212
6-5	Carbon isotope ratios of individual <i>n</i> -alkanes in a number of oils sourced from fluvio-deltaic (FD), lacustrine (LAC), marine deltaic (MD), and marine carbonate (MC) depositional environments. (Reprinted with permission from Murray et al. (1994). Copyright (1994) Elsevier Science.)	215
6-6	Carbon isotope ratios of individual <i>n</i> -alkanes in asphaltic bitumens from S. Australia (AGSO 934, 935), in a bitumen from N. Australia (AGSO 624), and in oils from Seram, Indonesia (AGSO 324) and Minas, Indonesia (AGSO 436). (Reprinted with permission from Dowling et al. (1995). Copyright (1995) Elsevier Science.)	216
6-7	Gas chromatograms and carbon isotope ratios of individual alkanes in oil from the feathers of birds killed in a major oil spill and from the oil that was the suspected source of the spill. (Reprinted with permission from Mansuy et al. (1997). Copyright (1997) American Chemical Society.)	217
6-8	Bulk carbon isotope ratios of oil from the Erika tanker spill (S1) off the Atlantic coast of France, 1999, and of oil residues collected from the northern (S2–S12) and southern (S13–S18) Atlantic coast. (Reprinted with permission from Mazeas and Budzinski (2002). Copyright (2002) American Chemical Society.)	218
6-9	Gas chromatograms and carbon isotope ratios of individual alkanes in beach tars from the Seychelles Islands. (Reprinted with permission from Philp (2002). Copyright (2002) Academic Press.)	219
6-10	Full scan GC/MS chromatograms of a sediment extract after sequential purification procedures. (A) Al/Si column chromatography (B) High performance liquid chromatography (C) Thin-layer chromatography. (Reprinted with permission from Kim et al. (2005). Copyright (2005) American Chemical Society.)	221
6-11	Carbon isotope ratios of individual naphthalenes and methylnaphthalenes in soils and sediments from McMurdo Station, Antarctica. (Reprinted with permission from Kim et al. (2005). Copyright (2005) American Chemical Society.)	222
6-12	Carbon isotope ratios of individual <i>n</i> -alkane/ <i>n</i> -alkene pairs generated by pyrolysing asphaltene fractions of biodegraded oils. These are compared with ratios of individual <i>n</i> -alkanes in the undergraded (initial) oils. (Reprinted with permission from Mansuy et al. (1997). Copyright (1997) American Chemical Society.)	223
7	Emerging CEN Methodology for Oil Spill Identification	
7-1	Decision/flowchart of the CEN oil spill identification methodology.	234
7-2	GC/FID chromatogram of a mildly weathered fuel oil displaying the dominating <i>n</i> -alkanes together with the acyclic isoprenoids, pristane and phytane (inserted).	235
7-3	Examples of weathering checks of gas oils by GC/FID. A. Simple overlaying of chromatograms. B. Bar chart comparison of normalized <i>n</i> -alkane distributions.	236
7-4	GC/MS ion fragmentograms of diagnostic alkylated PACs. 4A, methylphenanthrenes (<i>m/z</i> 192); 4B, methyl dibenzothiophenes (<i>m/z</i> 198); 4C, C2-phenanthrenes (<i>m/z</i> 206); 4D, C2-dibenzothiophenes (<i>m/z</i> 212);	

	4E, C3-phenanthrenes (m/z 220); 4F, C3-dibenzothiophenes (m/z 226); 4G, C4-phenanthrenes including retene (m/z 234); 4H, methylfluoranthenes/ methylpyrenes/benzofluorenes (m/z 216); 4I, C3-chrysenes (m/z 270).	239
7-5	Relative distribution of PAC homologues in two different IFO-180 bunker oils (from different refineries).	240
7-6	GC/MS ion fragmentogram of tri- and pentacyclic triterpanes (“hopanes”) recorded at m/z 191 in the SINTEF standard oil mixture.	241
7-7	GC/MS ion fragmentogram of tetracyclic rearranged (diasteranes) and regular 17α (H)-steranes, and 17β (H)-steranes recorded at m/z 217 and m/z 218, respectively, in the SINTEF standard oil mixture.	241
7-8	GC/MS ion fragmentogram of triaromatic steroids recorded at m/z 231 in the SINTEF standard oil mixture.	242
7-9	GC/MS ion fragmentogram of sesquiterpanes recorded at m/z 123 in the SINTEF standard oil mixture.	244
7-10	Protocol/decision chart for selection/elimination of diagnostic ratios.	247
7-11	Signal-to-noise calculation. The distance between the highest and lowest peaks of a part of the noise is used as N (see insert) and the peak height as S. In this example, S/N is 4.5.	249
7-12	PCA score plot of data (diagnostic ratios) from a recent mystery oil spill in the North Sea. Several possible source samples (production samples from nearby offshore fields) together with other production samples were included in the PCA.	250
7-13	GC/FID chromatogram of extracts of (A) candidate source sample (bilge), and (B) spill sample.	252
7-14	Weathering check. Overlay of GC/FID chromatograms of spill sample and candidate source sample.	253
8	Advantages of Quantitative Chemical Fingerprinting in Oil Spill Identification	
8-1	Comparison of two oils that exhibit qualitatively similar GC/FID patterns and selected extracted ion profiles (partial) that were interpreted to be “probable matches” by a laboratory conducting ASTM D3328 and D5739. Quantitative analysis showed some measurable differences existed, including higher concentrations of C2- and C3-dibenzothiophenes in oil 2 (see Fig. 8-2). *—internal standards.	262
8-2	Double-ratio plot showing the results of replicate analysis of the two oils shown in Figure 8-1. Despite qualitatively comparable D2 (m/z 212), D3, P2 (m/z 206), and P3 patterns (e.g., Fig. 8-1) the concentration of D2 and D3 were higher in Oil 2.	263
8-3	Examples of linear regression analysis as per the revised Nordtest oil spill identification method for the data shown in Table 8-4. (A) Example of “non-match” between spill oil and Source A based upon 98% confidence interval. (B) Example of “match” between spill oil and Source B based upon 98% confidence interval.	274
8-4	Revised Nordtest linear regression plots for ten diagnostic ratios (A-J) in the mystery spill oil and two candidate source oils from case study 1. (A) comparison of bilge oil with spilled oil resulting in a non-match, (B) comparison of HFO oil with spilled oil resulting in a non-match, (C) comparison of theoretical mixture of bilge and HFO with spilled oil resulting in a positive match.	278

- 8-5 Comparison of IFO and HFO source fuels oil fingerprints from a marine oil spill from case study 2. (A) GC/FID chromatogram for IFO, (B) GC/FID chromatogram for HFO, (C) PAH concentration histogram for IFO, (D) PAH concentration histogram for HFO, (E) Partial m/z 191 triterpane extracted ion profile for IFO, and (F) Partial m/z 191 triterpane extracted ion profile for HFO. H29 = Norhopane, H30 = Hopane * = internal standards. 280
- 8-6 Comparison of two field samples (mousse and tarball) associated with case study 2 and the vessel containing the fuels shown in Figure 8-5. (A) GC/FID chromatogram for the mousse sample (B) GC/FID chromatogram for the tarball sample, (C) PAH concentration histogram for the mousse sample, (D) PAH concentration histogram for the tarball sample, (E) Partial m/z 191 triterpane extracted ion profile for the mousse sample, and (F) Partial m/z 191 triterpane extracted ion profile for the tarball sample. H29 = Norhopane, H30 = Hopane * = internal standards. 281
- 8-7 Revised Nordtest linear regression plots for eleven diagnostic ratios (a-k) in the mousse sample and two candidate source oils (IFO and HFO) from case study 2. (A) Comparison of IFO with the mousse sample resulting in a non-match, (B) comparison of HFO oil with the mousse sample resulting in a non-match. 284
- 8-8 Revised Nordtest linear regression plots for eleven diagnostic ratios (a-k) in the tarball sample and two candidate source oils (IFO and HFO) from case study 2. (A) Comparison of IFO with the tarball sample resulting in a non-match, (B) comparison of HFO oil with the tarball sample resulting in a non-match. 285
- 8-9 Plot of the D2/P2 versus percent IFO and HFO from the grounded vessel. This equation was used with the D2/P2 ratios measured in the field samples to calculate the % IFO and % HFO in the field samples. These percentages were then used with the field sample quantitative data to calculate mixed oil source ratios that were used in the revised Nordtest method (e.g., Fig. 8-10). 286
- 8-10 Revised Nordtest linear regression plots for eleven diagnostic ratios (a-k) in the mousse (A) and tarball (B) samples versus the associated mixed source oils. The parity correlation was improved for both samples when the quantitatively determined mixed source ratios were used. Based on this analysis and considering the additional variability introduced by the mixing model but not reflected in the 98% confidence interval, a source match with the mixed oils is likely. 287
- 8-11 Revised Nordtest linear regression plots for eleven diagnostic ratios (a-k) in the tarball #2 sample versus the associated mixed source oil. No improvement in parity correlation was observed over the entire range of possible mixtures including the mixed source presented in this figure that was calculated based on the D2/P2 ratio in the sample. This data combined with GC/FID, PAH and biomarker signatures is conclusive evidence that this sample is not related to the oil spill discussed in case #2. 289
- 9 A Multivariate Approach to Oil Hydrocarbon Fingerprinting and Spill Source Identification
- 9-1 Illustration of the research strategy behind the IMOF methodology, which is based on four steps: chemical analysis, data preprocessing, multivariate data analysis and data evaluation. The aim of the work in our research groups has

	been to develop rapid, reliable and objective tools for comprehensive analysis and characterization of complex oil hydrocarbon mixtures using the IMOF methodology.	297
9-2	Fluorescence excitation-emission scans of an HFO. The vertical axis and scale bar show the fluorescence intensity. Modified from Christensen et al. (2005b). See color plate.	299
9-3	Partial GC-MS chromatogram of m/z 217, which contains tricyclic steranes (eluting between 26 and 34 min) and tetracyclic steranes (eluting between 35 and 42 min).	301
9-4	First derivative of a section of m/z 217 for five reference oils and five source oils: (a) before warping and (b) after warping using COW with segment length of 175 data points and a slack of three points. Modified from Christensen et al. (2005d).	303
9-5	Preprocessed fluorescence EEM of the same oil sample as shown in Figure 9-2 after blank subtraction, insertion of missing values, and a small triangle of zeros and excitation/emission correction. The vertical axis and scale bar show the fluorescence intensity. Modified from Christensen et al. (2005b). See color plate.	308
9-6	(a) Weights ($RSD_{\bar{A}}^{-1}$) used for WLS-PCA and (b) the aligned and normalized mean combined chromatogram of unweathered Brent crude oil. The mean was calculated from four replicate samples. Modified from Christensen et al. (2005a).	314
9-7	Score plot (PV2 versus PC4) using WLS-PCA on sections of preprocessed partial GC-MS/SIM chromatograms of m/z 217. The PCA model was calculated from the calibration set (61×1231), whereas a reference set (18×1231) of replicate references and test set (22×1231) were calculated by projecting the data onto the loadings. The test set was comprised of 16 <i>Baltic Carrier</i> oil spill samples and two spill samples from a Round-Robin exercise analyzed in triplicate (Spill I and Spill II) (Faksness et al., 2002).	316
9-8	Integrated mean-chromatogram (dotted line) and integrated PC4 loadings (solid line) for WLS-PCA in the oil hydrocarbon fingerprinting study in Christensen et al. (2005d). The $\alpha\alpha$ -steranes ($\alpha\alpha$) and $\beta\beta$ -steranes ($\beta\beta$) are marked in the plot.	316
9-9	PARAFAC score plot. Factor 3 versus Factor 5. Symbols for LFOs, HFOs, lubes, crude oils, unknown oil samples, replicate reference oils, and the triplicate spill and ship samples are explained in the legend. The 17 replicate references are circled, and arrows mark the position of spill and <i>Baltic Carrier</i> ship samples.	318
9-10	Comparison of PARAFAC excitation and emission loading spectra of selected PAHs. Loadings and spectra of individual PAHs have been normalized to ease visual comparison. Furthermore, excitation and emission loadings are shown together in the plots, where excitation loadings appear to the left of the corresponding emission loadings.	318
9-11	Flowchart for the IMOF methodology including the individual oil hydrocarbon fingerprinting methods developed by Christensen et al. (2004, 2005b, 2005c, 2005d).	320
10	Chemical Heterogeneity of Marine Heavy Fuel Oils	
10-1	Gas chromatographic “fingerprints” of six IFO-380 HFOs, demonstrating the significant variability in chemical compositions in the sample fuel class. * — internal standards.	334

- 10-2 Ternary diagram depicting the variability in the bulk hydrocarbon composition among 71 worldwide IFO 380 heavy fuel oil samples. Weight percentages based upon quantitative GC/FID analysis and gravimetric analysis. 335
- 10-3 Ternary diagrams showing the distribution of C₂₇, C₂₈, and C₂₉ regular steranes (5 α , 14 β , 17 β (H) 20S+20R) calculated from absolute concentrations determined from the *m/z* 218 mass chromatogram for 71 IFO 380 residual fuel oils. 337
- 10-4 Cross-plots of (A) triterpane-based and (B) sterane-based thermal maturity parameters for 71 IFO 380 fuels. Dashed lines show typical maximum (equilibrium) values for crude oils (from Peters et al., 2005). Triterpane ratios both measured on the *m/z* 191 mass chromatograms. 14 α (H), 17 α (H) C₂₉ steranes measured on the *m/z* 217 mass chromatogram and 14 β (H), 17 β (H) C₂₉ steranes measured on the *m/z* 218 mass chromatogram. 337
- 10-5 Average (± 1 standard deviation) concentration of select PAH compounds in 71 worldwide IFO 380 HFOs. See Table 10-3 for compound abbreviations. Inset shows expanded view of 4- to 6-ring PAH. 340
- 10-6 Percentage C₀–C₄ naphthalenes of the total PAH in 71 worldwide IFO 380 samples. Total PAH represents sum of 54 analytes from Table 10-3. 340
- 10-7 Histograms comparing distribution and concentration of PAH in (A) low aromatic IFO-380 and (B) high aromatic IFO-380. Gas chromatogram for (C) high aromatic, cracked gas oil blending stock, and (D) a high aromatic IFO-380 containing a prominent cracked gas oil blending stock. 341
- 10-8 Cross-plots of (A) the range in of the diagnostic source ratios D2/P2 and D3/P3 and (B) the relationship between D2/P2 and total PAH in 71 worldwide IFO 380 heavy fuel oil samples. 343
- 10-9 Structures of (A) linear 3-ring anthracene molecule and (B) nonlinear 3-ring phenanthrene molecule. 343
- 10-10 GC/FID chromatograms for (A) an unweathered IFO 380 fuel and (B) moderately evaporated crude oil and partial *m/z* 192 mass chromatograms for (C) the unweathered IFO 380 shown in (A) and (D) the moderately evaporated crude oil shown in (B). N — alkyl-naphthalenes, P — alkyl-phenanthrenes; MP — methylphenanthrene, MA — methylanthracene. 344
- 11 Biodegradation of Oil Hydrocarbons and Its Implications for Source Identification
- 11-1 Typical reactions catalyzed by hydro-carbon monooxygenases. Octane monooxygenase of *Pseudomonas putida* catalyzes the oxidation of octane to octanol (van Beilen et al., 1994), toluene-3-monooxygenase of *Ralstonia pickettii* catalyzes the oxidation of toluene to 3-cresol (Tao et al., 2004), the cytochrome P450 of *Mycobacterium vanbaalenii* apparently catalyzes the oxidation of benzo[a]pyrene to benzo[a]pyrene-11,12-epoxide (Moody et al., 2004), and the cyclohexane monooxygenase of *Brachymonas petroleovorans* oxidizes cyclohexane to cyclohexanol (Brzostowicz et al., 2005). Other enzymes catalyze the subterminal oxidation of alkanes (Ludwig et al., 1995), and the oxidation of the 2- or 3-position of toluene (Yeager et al., 1999; Tao et al., 2004). 351
- 11-2 Typical reactions catalyzed by hydro-carbon dioxygenases. The alkane dioxygenase of an *Acinetobacter* apparently converts alkanes to alkanehydroperoxides (Maeng et al., 1996). Naphthalene dioxygenase of *Pseudomonas putida* oxidizes naphthalene to *cis*-naphthalene-1,2-dihydrodiol (Karlsson et al., 2003). 352

- 11-3 The opening of aromatic rings by extradiol and intradiol dioxygenases. Examples include the extradiol protocatechuate 4,5-dioxygenase of *Sphingomonas paucimobilis* (Sugimoto et al., 1999) and the intradiol protocatechuate 3, 4-dioxygenase of *Acinetobacter* (Vetting et al., 2000). 353
- 11-4 The biodegradation of trimethyl, methyl-ethyl, propyl, and isopropyl benzenes by four organisms. *Pseudomonas putida* F1 degrades toluene with a dioxygenase (Cho et al., 2000), while *Ps. putida* mt-2 (Bühler et al., 2000), and *Ps. mendocina* KR (Tao et al., 2004) use monooxygenases, directed at the methyl group and the *para*-position, respectively. *Sphingomonas yanoikuyae* B1 possesses a biphenyl dioxygenase and a xylene monooxygenase (Kim and Zylstra, 1999), and it is certainly possible that all strains contain other hydrocarbon-oxidizing enzymes. These experiments (Prince, V.L., Zylstra, G.J., and Prince, R.C., unpublished) used cells initially grown with vapor phase toluene that were washed and resuspended in minimal medium with 1 µl of gasoline in 10 ml of culture in a 40-ml vial and incubated for 90–113 hr. The traces are for the *m/z* = 105 ion. 354
- 11-5 Anaerobic hydrocarbon activation by fumarate addition. The denitrifying bacterium HxN1 catalyzes the production of 1-methylpentylsuccinate from hexane (Rabus et al., 2001) and Biegert et al. (1996) were the first to demonstrate the production of benzylsuccinate under anaerobic conditions with *Thauera aromatica*. Similar reactions have been seen under a variety of conditions by a diverse array of organisms (Rabus, 2005). 355
- 11-6 Anaerobic hydrocarbon activation by carboxylation and dehydration. Carboxylation at the 2-carbon of hexadecane by the sulfate-reducing strain AK-01 was reported by So and Young (1999). Carboxylation at the 3-position was subsequently reported for strain Hxd3 (So et al., 2003). Carboxylation of naphthalene by a sulfate-reducing enrichment culture was reported by Zhang and Young (1997). Recent work on benzene degradation (Chakraborty and Coates, 2005; Ulrich et al., 2005) suggests that a transient hydroxylation precedes carboxylation. While it seems clear that labeled bicarbonate is incorporated into the hydrocarbons, the enzymes responsible have not been characterized. In contrast, ethylbenzene dehydrogenase from *Azoarcus* has been well-characterized (Johnson et al., 2001; Kniemeyer and Heider, 2001); it contains the molybdopterin cofactor. 356
- 11-7 Anaerobic degradation of hexadecane under methanogenic conditions. A consortium of syntrophic eubacteria and archaea can degrade hexadecane to CH₄, CO₂, and H₂S. [Figure adapted from Parkes (1999) and Zengler et al. (1999)]. 357
- 11-8 Bulk oil properties for a suite of oils (La Luna source) from Eastern Venezuela. Sulfur, nitrogen, nickel, and vanadium increase proportionally with decreasing API gravity. The degree of biodegradation is indicated by the numerical ranking of the Biomarker Biodegradation Scale (Peters et al., 2005) and by descriptive terms used by Wenger et al. (2002). 358
- 11-9 Gas chromatograms for a suite of selected crude oils from Africa. The oils were generated and expelled from the same source rock under comparable thermal conditions, but indicate quite different amounts of *in situ* biodegradation. Shown are reservoir temperature, API gravity, viscosity, pristane/phytane (Pr/Ph) and *n*C₁₇/pristane (*n*C₁₇/Pr) ratios, and mass percentage lost calculated using hopane as a conserved internal standard.

	Number labels indicate <i>n</i> -alkane carbon number [modified from Wenger et al. (2002)].	359
11-10	The extent of biodegradation of mature crude oil can be ranked on a scale of 1–10 based on differing resistance of compound classes to microbial attack. Biodegradation is quasi-sequential because some of the more labile compounds in the more resistant compound classes can be attacked prior to complete destruction of less resistant classes [from Peters et al. (2005)]. Arrows indicate where compound classes are first altered (dashed lines), substantially depleted (solid gray), and completely eliminated (black). Sequence of alteration of alkylated PAHs is based on work by Fisher et al. (1998) and Trolio et al. (1999). Qualitative descriptions of the degree of biodegradation from Wenger et al. (2002) reflect changes in oil quality.	361
11-11	Differential biodegradation within compound classes. Modified from Peters et al. (2005).	362
11-12	25-norhopanes have the same optical configuration as hopanes. They differ only by the removal of the methyl group attached to the C-10 carbon.	363
11-13	Mass chromatograms of hopanes (<i>m/z</i> 191) and norhopanes (<i>m/z</i> 177) from Eastern Venezuela. Ion traces are scaled proportional to their relative response. The 25-norhopanes (indicated by D-carbon number) are believed to originate by loss of a methyl group from C-10 in hopanes. Thus, the single epimer of C ₃₀ 17 α ,21 β (H)-hopane (top) has been partially altered to C ₂₉ 25-nor-17 α -hopane (bottom), while each of the C ₃₁ –C ₃₅ 17 α -hopane (22S + 22R) epimers correspond to two C ₃₀ to C ₃₄ 25-norhopane epimers. Vertical lines indicate some peaks that yield both <i>m/z</i> 191 and 177 ions. Modified from Peters et al. (2005).	363
12	Identification of Hydrocarbons in Biological Samples for Source Identification	
12-1	Typical weathering sequence of PAH in Alaska North Slope crude oil. N = naphthalene, F = fluorene, D = dibenzothiophene, P = phenanthrene/anthracene, C = chrysene; numbers indicate substituent alkyl carbon atoms. Weathering increases from top panel (unweathered oil) to bottom panel (moderately weathered oil). Abscissa: proportion of total PAH.	385
12-2	Relative composition of aqueous PAH dissolved from initially unweathered Alaska North Slope oil. Abbreviations and axes as in Figure 12-1.	386
12-3	Relative abundance of PAH and phytane in (A) <i>Exxon Valdez</i> oil mousse 11 days following the spill, and (B) in mussels collected six weeks later. Note the PAH weathering evident in the mussels compared with the oil (compare with Figure 12-1), and the prominent phytane in both, indicating ingestion of whole weathered oil droplets by the mussels. Abbreviations: Naph = naphthalene, Menaph = methylnaphthalene, Acenthy = acenaphthylene, Acenthe = acenaphthene, Fluor = fluorene, Dithio = dibenzothiophene, Phenanth = phenanthrene, Anthra = anthracene, Fluorant = fluoranthene, Benanth = benzo[<i>a</i>]anthracene, Chrys = chrysene, Benzobfl = benzo[<i>b</i>]fluoranthene, Benzokfl = benzo[<i>k</i>]fluoranthene, Benepyr = benzo[<i>e</i>]pyrene, Beneapyr = benzo[<i>a</i>]pyrene, Indeno = indeno[1,2,3- <i>c,d</i>]pyrene, Dibenz = dibenzo[<i>a,h</i>]anthracene, Benzop = benzo[<i>g,h,i</i>]perylene (from Short and Harris, 1996b).	387
12-4	Cytochrome P450 1A induction pathway (from Whitlock, 1993; see text in section 12.3.1 for abbreviation).	389

12-5	Gill epithelial cells under normal conditions (left), and under CYP 1A induction (right) (from Malins et al., 2004).	389
12-6	Catalysis of 7-ethoxyresorufin to resorufin by cytochrome P450 1A (CYP 1A). Resorufin is fluorescent, and its production rate is the basis for the EROD assay for CYP 1A activity.	390
12-7	Some effects of oil exposure in herring larvae (photos courtesy of NOAA).	395
13	Trajectory Modeling of Marine Oil Spills	
13-1	Comparison of a trajectory forecast with oil representing 100 (a) and 1000 (b) particles or Lagrangian elements (LEs).	409
13-2	A cross-sectional view of an idealized oil spill in the sea. The arrows below the oil streaks represent the motion of these vortices. The oil is shown collecting in areas with converging surface currents.	414
14	Oil Spill Remote Sensing: A Forensics Approach	
14-1	Spectra of three fuels with similar physical properties showing the spectral differences in them using a fluorosensor.	425
14-2	The return signals from the LURSOT thickness sensor showing a measurement of 9 mm from a light crude oil. Measurement was taken 61 m above ground with aircraft at 200 km/hr.	435
15	Advances in Forensic Techniques for Petroleum Hydrocarbons: The Exxon Valdez Experience	
15-1	PAH compositions of major hydrocarbon source inputs to the marine environment of PWS. A PAH analyte may be a single compound, such as C ₀ -phenanthrene, or the sum of a group of isomers, such as C ₁ -phenanthrene, which has four isomers. N1, N2, N3, N4 = C ₁ , C ₂ , C ₃ , C ₄ -naphthalenes; ACL = acenaphthylene; ACE = acenaphthene; BPH = biphenyl; F, F2, F2, F3 = C ₀ , C ₁ , C ₂ , C ₃ -fluorenes; P, P1, P2, P3, P4 = C ₀ , C ₁ , C ₂ , C ₃ , C ₄ -phenanthrenes/anthracenes; D, D1, D2, D3 = C ₀ , C ₁ , C ₂ , C ₃ -dibenzothiophenes; FL = fluoranthene; PY = pyrene; FP1 = C ₁ -fluoranthenes/pyrenes; BaA = benzo(a)anthracene; C, C1, C2, C3, C4 = C ₀ , C ₁ , C ₂ , C ₃ , C ₄ -chrysenes; BbF = benzo(b)fluoranthene; BkF = benzo(k)fluoranthene; BeP = benzo(e)pyrene; BaP = benzo(a)pyrene; Per = perylene; IDP = indeno(1,2,3-cd)pyrene; DBA = dibenzo(a,h)anthracene; BgP = benzo(g,h,i) perylene.	452
15-2	PAH distributions, TPAH and 3–4-ring PAH concentrations, major fraction compositions, and weathering parameters for EVC and four variably weathered PWS shoreline oil samples. The weathering parameter, ω, calculated from the distributions of 14 PAH analytes (Short and Heintz, 1997) becomes indeterminate (IND) at advanced weathering states when one or more analytes are not detected.	454
15-3	Pyrogenic index FP/(FP + C24Ph) vs. the ratio of low-molecular-weight (2–3-ring PAH) to total PAH, LPAH/TPAH, for naturally and artificially weathered shoreline oils. LPAH/TPAH decreases with increased weathering. 1989–1990 bioremediation suite oils are from Prince et al. (1994). EM shoreline oil weathering suite from samples collected in 2001–2002. NOAA EVTHD oils from the 2003 version of Exxon Valdez Trustees Oil Database on	

- NOAA-NMFS, Auke Bay Laboratory webpage (www.afsc.noaa.gov/abl/oilspill). These include oils that were weathered in the laboratory. 458
- 15-4 PAH and triaromatic steroid (TAS) compositions (C_2 -phenanthrene-normalized) for (A) PWS prespill segment of a benthic core, (B–J) potential source inputs to the natural background. Abbreviations: Np = naphthalenes, Fl = fluorenes, Ph = phenanthrenes, Db = dibenzothiophenes, Ch = chrysenes, Per = perylene, N0 = C_0 -naphthalene, N1 = C_1 -naphthalene, P0 = C_0 -phenanthrene, P1 = C_1 -phenanthrene, P2 = C_2 -phenanthrene, Db2 = C_2 -dibenzothiophene. Adapted with permission from Boehm et al. (2001). Copyright 2001, American Chemical Society. 460
- 15-5 Triterpane and sterane compositions (C_2 -phenanthrene-normalized) for (A) PWS benthic core, (B–J) potential source inputs to the natural background. Abbreviations: Ts = $17\alpha(H)$ -22,29,30-trisnorhopane, Tm = $18\alpha(H)$ -22,29,30-trisnorhopane, 28B = $C_{28,30}$ -bisnorhopane, Dpl(ROM) = diplotene, recent organic matter (the Copper River is a major source), 29Nor = norhopane, Hop = C_{30} -hopane, Diaster = diasteranes. Adapted with permission from Boehm et al. (2001). Copyright 2001, American Chemical Society. 461
- 15-6 C_2 -Dibenzothiophenes *versus* C_2 -phenanthrenes (D2/P2) relationship of weathered EVC, diesel refined from ANS crude feed stock at a Kenai Refinery, PWS regional background hydrocarbons from deep subtidal cores, and Katalla and Cook Inlet crude oils. 462
- 15-7 Sediment PAH compositions for a profile from the mid-tide zone to 100-m water depths at a location containing buried EVC residues on the beach, Northwest Bay, Eleanor Island. (Data from S. Rice, personal communication, 2003). 463
- 15-8 Factor score plot resulting from the PCA analysis of PAH in shoreline and benthic sediments in PWS. Modified after Burns et al. (1997). 465
- 15-9 Source apportionments for GOA and prespill PWS benthic sediments. KB = Katalla Beach; MGF = Malaspina Glacier Flour. Reprinted with permission from Boehm et al. (2001). Copyright 2001, American Chemical Society. 467
- 15-10 1999–2000 embayment benthic sediment sampling program with site locations, oiling category, representative sediment PAH profiles, and the results of the CLS source allocations, where ANS = Alaska North Slope oil, Mon = Monterey petroleum, PYR = pyrogenic PAH, and BKG = regional petrogenic background. 470
- 15-11 PAH and alkane compositions of bald eagle eggshells. (A) Weathered EVC on eggshell collected at Italian Bay on June 8, 1989, EVTHD ID #23363. (B) Relatively fresh diesel on eggshell collected from Hells Hole in eastern PWS outside the spill zone on July 18, 1990, EVTHD ID #20029. 472
- 15-12 PAH and alkane compositions for sea otter tissues. (A) Heavily weathered EVOS, sea otter skin collected April 12, 1989, EVTHD ID #27757, site not documented. (B) Relatively fresh diesel, sea otter fur collected August 1, 1990, site not documented, EVTHD ID #22851. (C) Procedural artifact, sea otter blood, Ogden Passage, SE Alaska, EVTHD ID #20044. 473
- 15-13 PAH compositions for Harlequin duck stomach contents (A–B) and corresponding liver tissue (C–D) from the same bird collected by State of Alaska investigators. Adapted, with permission, from *STP 1219 Exxon Valdez Oil Spill: Fate and Effects in Alaskan Water*, copyright ASTM International, 100 Barr Harbor Drive, West Conshohocken, PA 19428. 474

15-14	PAH compositions of tissue exposed to weathered oil in column generator experiments conducted by NOAA/ABL, Juneau, AK. A. Pink salmon eggs after 36 days of exposure to artificially weathered EVC at a gravel loading of 281 ppm oil showing characteristic WSF composition. B. Herring eggs showing mixed oil + WSF composition.	476
15-15	Composition and concentration of PAH in mussel tissues from NOAA oiled mussel bed sites compared with the PAH composition and concentration of underlying sediments for samples collected in 2001. The data are presented as the geometric means of the analyte distributions of replicate samples at each site. Note that the mussel TPAH concentrations are factors of 3 to 300 lower than the TPAH concentrations in the corresponding sediments.	478
15-16	Mean sediment TPAH versus composite clam TPAH for 16 oiled, 3 reference, and 4 HA sites in western PWS. UIZ = upper intertidal zone (+2 m mllw), MIZ = middle intertidal zone (+1–+2 m mllw), LIZ = lower intertidal zone (0–+1 m mllw).	479
15-17	PAH composition of composite clam tissues from two sea otter foraging sites that had been oiled in 1989.	480
15-18	Correlation of rockfish liver ethoxyresorufin <i>o</i> -deethylase (EROD) activity, a measure of the level of CYP 1A induction, with benthic sediment pyrogenic index (FP)/(FP + C ₂₄ Ph). Solid symbols = sites oiled in 1989. Open symbols = unoiled reference sites.	482
16	Case Study: Oil Spills in the Strait of Malacca, Malaysia	
16-1	Map of Peninsular Malaysia showing the Straits of Malacca.	490
16-2	Oil spill incidents in the Straits of Malacca (Malaysian Marine Department, 2004).	491
16-3	Sampling stations for tar ball samples (solid diamond) in Peninsular Malaysia.	493
16-4	Sampling stations for sediment in Peninsular Malaysia. Solid triangles represent river stations and solid circles represents inshore stations.	494
16-5	Cross-plots for source identification of oils. The circles indicating both categories were established through the analysis of crude oils in Zakaria et al. (2000).	501
17	Evaluation of Hydrocarbon Sources in Guanabara Bay, Brazil	
17-1	Social, economical, and physical characteristics of Guanabara Bay.	507
17-2	Oil slick after January 2000 oil spill accident in Guanabara Bay.	508
17-3	Geographical location of the 21 sediment stations in Guanabara Bay sampled in 2000 and 2003.	509
17-4	Predominance of alkylated PAH for intertidal samples from Guanabara Bay 2000 survey in relation to 2003.	517
17-5	Predominance of four- and five-ring PAH for subtidal samples from Guanabara Bay for 2000 and 2003 campaigns. (a) Inside the influence of the oil spill slick; (b) outside the influence of the oil spill slick.	518
17-6	PAH cross-plot of phenanthrene/anthracene versus fluoranthene/pyrene for Guanabara Bay sediments and MF 380, Arabian oil (AL), light oil (Ar), and diesel oil (DM).	520
17-7	PAH cross-plot of anthracene/(anthracene + phenanthrenes) versus fluoranthene/(fluoranthene + pyrene) for Guanabara Bay sediments from	

	campaigns 2000 and 2003 and MF 380, Arabian oil (AL), light oil (Ar), and diesel oil (DM).	520
17-8	PAH cross-plot of indeno1,2,3-(cd)pyrene/(indeno1,2,3-(cd)pyrene + benzo(ghi)perylene) versus fluoranthene/(fluoranthene + pyrene) for Guanabara Bay sediments from campaigns 2000 and 2003 and MF 380, Arabian oil (AL), light oil (Ar), and diesel oil (DM).	521
17-9	PAH cross-plot of fluoranthene/(fluoranthene + pyrene) versus phenanthrene + anthracene/(phenanthrene + anthracene + C1phenanthrenes) for Guanabara Bay sediments from campaigns 2000 and 2003 and MF 380, Arabian oil (AL), light oil (Ar), and diesel oil (DM). Interpretive guidelines based upon Yunker et al. (2000).	522
17-10	Plot of the relative ratios Σ (other 3–6-ring PAH)/ Σ (5 alkylated PAH series) versus phenanthrene/anthracene for Guanabara Bay sediments from the campaigns 2000 and 2003 and MF 380, Arabian oil (AL), light oil (Ar), and diesel oil (DM). Interpretive guidelines based upon Wang et al. (1999b).	523
17-11	Plot of the relative ratios Σ (other 3–6-ring PAH)/ Σ (5 alkylated PAH series) versus fluoranthene/(fluoranthene + pyrene) for Guanabara Bay sediments from the campaigns 2000 and 2003 and MF 380, Arabian oil (AL), light oil (Ar), and diesel oil (DM). Interpretive guidelines from Wang et al. (2002b).	524
17-12	Relative perylene abundance for Guanabara Bay sediments from the campaigns 2000 and 2003.	525
17-13	PCA projections of PAH variables and sediment samples from campaign 2000 and MF 380, Arabian oil (AL), light oil (Ar), and diesel oil (DM).	526
17-14	PCA projections of PAH variables and sediment samples from campaign 2003 and MF 380, Arabian oil (AL), light oil (Ar), and diesel oil (DM).	526
17-15	Terpane and sterane partial mass chromatograms of the spilled marine heavy fuel oil, January 2000.	527
17-16	A plot of C27/C29-steranes 5 α ,14 β ,17 β (H),20S against C29/C27-steranes 5 α ,14 α ,17 α (H),20S showing the similarity between most of the samples and the correlation between both samples T24 collected in campaigns 2000 and 2003.	531
17-17	A plot of C24-tetracyclic/C26-tricyclic terpanes against C29/(C29 + C30-hopanes) highlighting the correlation between the marine fuel and samples T7 and T9 from the campaign 2000.	531
17-18	A plot of Ts/(Ts + Tm) against 30-norhomohopane/(30-norhomohopane + hopane) showing the correlation between samples and a marine carbonate-sourced crude oil.	531
17-19	A plot of C27/C29-steranes 5 α ,14 β ,17 β (H),20S against C29/C27-steranes 5 α ,14 α ,17 α (H),20S showing the correlation between most of the samples and a marine carbonate-sourced crude oil.	532
17-20	A plot of C27/C29-steranes 5 α ,14 β ,17 β (H),20S against C28/C29-steranes 5 α ,14 β ,17 β (H),20S showing the correlation between the sediment samples and a marine carbonate-sourced crude oil.	532