

Index

A

- Abietane, 14
- Aboveground storage tanks (ASTs), 31
- Accidental oil spills, 1–2
- Acenaphthalene, 514
- Acenaphthene, 8, 267, 339, 514
- Acenaphthylene, 5, 8, 267, 339
- Acetate, 356–357
- Acetogens, 357
- Acid rain gases, 150
- Acids, 360
- Acinetobacter*, 352
- Acoustic systems, 435
- Acyclic acids, 361
- Acyclic isoprenoids, 14
- Adamantanes, 114–117
 - diagnostic ratios, 127
 - empirical formula, 78
 - mass spectra of, 88
 - target ions, 78
- Aegean Sea* oil spill, 162
- Aerobic biodegradation, 351–355
- Aerobic respiration, 366, 367–368
- Aerosol, 495
- Air monitoring, 59–60
- Air-oil interface, 429
- Aircraft fuel, 2
- Alaska North Slope (ANS) crude oil, 9, 16–17, 93–99, 111–113
- Alberta Sweet Mix Blend (ASMB) crude oil, 9, 161, 434
- Alcanivorax*, 353
- Aldehydes, 155
- Algal debris, 38–39
- Alicyclic compounds, 356
- Aliphatic hydrocarbons, 14
- Alkane alcohols, 352
- Alkane carboxylase, 356
- Alkane dioxygenase, 352–353
- Alkanes, 14, 300, 353, 361, 496–497
- Alkenes, 352
- Alkylated benzenes, 362
- Alkylated benzothiopenes, 152
- Alkylated naphthalenes, 471
- Alkylbenzenes, 132, 176–177
- Alkylbiphenyls, 362
- Alkylcyclohexanes, 24–26, 361
- Alkyldiphenylmethanes, 362
- Alkyltoluenes, 300
- Alpha hydrogens, 80
- Ameiurus nebulosus*, 394–396
- Amoco Cadiz* oil spill, 5, 13
- Anaerobic biodegradation, 355–357
- Anaerobic respiration, 368–369
- Anthracenes, 5, 8, 267, 339, 514
- Anthracite, 40
- Anthropogenic background hydrocarbons, 41–43
- API gravity, 350, 354, 359–360
- Arabian Light crude oil, 93–99, 111–113
- Arcachon Bay, 516
- Archaea, 198
- Argo Merchant*, 406, 407
- Aromatic dioxygenases, 353
- Aromatic hydrocarbons, 14–15
- Aromatic steranes, 77, 104–109
- Aromatic steroids, 240–244, 361, 362
- Arrow* oil spill, 133, 449
- Artemia*, 506
- Aryl hydrocarbon nuclear translocator protein (ARNT), 388
- Aryl hydrocarbon receptor (AhR), 388–389
- Asphaltenes, 13, 211
- Asphaltic bitumens, 216
- Asphalts, 345–346, 494
- ASTM 5623-94, 157
- ASTM D3328, 5, 259–262, 330
- ASTM D5739, 5, 259–262, 330–331
- Asymmetric carbons, 80–81
- Atmospheric deposition, 2
- Atmospheric gas oils, 22
- Atomic emission detection (AED), 157
- Automotive gasoline, 22
- Aviation gasoline, 22, 350
- Azoarcus*, 355–356

B

- Baccharane, 90
- Bacillus*, 364
- Background chemicals, 36–37
 - anthropogenic, 41–43
 - river runoff, 42–43
 - urban, 42–43
 - chemical fingerprinting of, 37

- Background chemicals (*Continued*)
- naturally-occurring, 37–41
 - algal debris, 38–39
 - oil seeps, 41
 - particulate coal, 39–41
 - vascular plants, 38–39
 - wood charcoal, 39–41
 - statistical analysis of, 37
 - types of, 36
- Bacteria, 350
- Bacteriocides, 364
- Baffin Island oil spill, 354
- Baltic Carrier* oil spill, 295, 315, 319
- Barge vessel spills, 2
- Base peak, 84
- Baselines, 303–304
- Battelle Memorial Institute, 230
- Beach tars, 215–216
- Beaufort Sea, 437
- Benchtop quadropole GC/MS, 84
- Benzenes, 5, 354–355
- Benzo ring, opening of, 155
- Benzoanthracene, 5, 267
- Benzofluoranthenes, 5, 8, 267, 339, 514
- Benzofluorenes, 237–239
- Benzopyrenes, 5, 8, 267, 339, 514
- Benzopyrenes, 5, 8, 267, 339
- Benzothiophenes, 27, 149–150, 152, 300
- Benzylsuccinate synthase, 355
- Beta*-carotane, 124
- Beta* hydrogens, 80
- Bicadinanes, 123–124
- Bicyclic terpanes, 361
- Bilge water, 28
- Bioavailability assessment, 477–482
 - biomarkers, 481–483
 - passive sampling of PAH, 481
 - uptake of PAH in biota, 477–482
- Biodegradation, 349–370. *See also* Weathering
 - aerobic, 351–355
 - anaerobic, 355–357
 - biochemistry of, 349–350
 - and biomarker fingerprinting, 131
 - of biomarkers, 360–361
 - and chemical fingerprinting, 32–34
 - and chemical transformation of crude oil, 153
 - limiting factors, 362–364
 - access of microbes to oil, 363–364
 - lack of bacteriocides, 364
 - nutrients, 363
 - presence of microorganisms, 364
 - temperature, 364
 - terminal electron acceptors, 362–363
 - water, 364
 - microbial ecology of, 365–369
 - deep subsurface ecology, 367–369
 - aerobic respiration, 367–368
 - anaerobic respiration, 368–369
 - microbial communities, 365–367
 - in reservoirs, 357–358
 - subsurface, 357–362
 - susceptibility to, 362
- Biogenic hydrocarbons, 350, 451, 453
- Biomarker fingerprinting, 73–74
 - analysis methods for, 81–83
 - application to oil spill studies, 117–121
 - capillary GC/MS analysis in, 83–86
 - effects of weathering on, 130–138
 - biodegradation, 131
 - dissolution, 131
 - emulsification, 130
 - evaporation, 130
 - natural dispersion, 130–131
 - oil-mineral aggregation, 132
 - photooxidation, 131–132
 - sedimentation, 132
 - internal standards, 83
 - source identification of spills, 134–138
 - characterization of sesquiterpanes, 135
 - evaluation of alkylated PAHs, 135–138
 - evaluation of pentacyclic terpanes and steranes, 135–138
 - product type-screening, 134–135
 - surrogates, 83
- Biomarkers, 9–10
 - biodegradation of, 133–134, 360–361
 - compound classes, 16, 74–77
 - cyclic terpenoids, 75–77
 - isoprenoids, 75
 - cross-plots of, 128–130
 - for oil and petroleum products, 129–130
 - for spill source identification, 129
 - in crude oils, 93–99
 - definition of, 15
 - diagnostic indices, 269–272
 - diagnostic ratios of, 126–128, 239–240, 511, 514
 - in heavy fuel oils (HFO), 336–338
 - identification of, 90
 - labeling and nomenclature of, 77–81
 - asymmetric carbons, 80–81
 - R and S stereoisomers, 81
 - stereoisomers, 79–80
 - in lubricating oils, 101–103
 - mass fragments, 300

- mass spectra of, 86–90
- in oil bioavailability assessment, 481–483
- in oil fractions, 104
- overview, 73–74
- in petroleum products, 99–101
- in qualitative fingerprinting, 264–265
- source-specific, 121–125
- weathered percentages, 134
- Bioresistance, 360
- BIOS oil spill, 133
- Biosphere, 349
- Biphenyl dioxygenase, 354
- Biphenyls, 8, 178–180, 267, 300, 339
- Biphytane, 198
- Birds, effects of oil spills on, 216–217
- Bishomohopanes, 242, 267
- Bisnorhopane, 242, 267
- Bitumens, 215–216
- Bituminous coal, 40
- Blue-sac disease, 394–395
- Boil range indices, 271
- Boiling points, 351
- Bonn Agreement, 229
- Boscan crude oil, 96–99
- Botryococcane, 75, 123, 215
- Botryococcus braunii*, 121, 125
- Bouchard 120*, 191
- Bouchard 65*, 187–189
- BP American Trader*, 212–213
- Brachymonas petroleovorans*, 351
- Branched alkanes, 362
- Branched hydrocarbons, 352
- Brevibacterium*, 352
- Brisbane River, 516
- Bromochloriodomethane, 81
- Bulk isotope ratios, 210–213
- Bunker C fuel oil, 101, 191–195, 329
- Bunker fuel, 161, 327, 329
- Buzzards Bay oil spill, 191–195
- C**
- C-band radar, 427
- Cadalene, 267
- Cahn-Ingold-Prelog convention, 81
- Calcite, 196
- Calcium, 329
- California crude oil, 96–99, 111–113
- Cameras, 420–422
- Capillary columns, 6
- Carbon dioxide, 369
- Carbon isotope ratios, 209–212
- Carbon isotopes, 208
- Carbon range analysis, 332
- Carboxylated pyrrols, 361
- Carboxylation, 356
- Carboxylic acids, 155
- Cargo washings, 2
- Carotane, 124
- Casco Bay, 516
- Catabolic degradation, 387–393
- Catalytic cracking, 24
- Catechols, 352
- CEN oil spill identification methodology, 229–254.
 - See also* Oil hydrocarbon fingerprinting
 - case study, 251–254
 - final evaluation and conclusions, 250–251
 - overview, 230–231
 - scope of, 231
 - strategies, 231–233
 - tiered levels of analysis and data treatment in, 233–251
 - decision chart, 233–234
 - GC/FID screening (level 1), 235–237
 - GC/MS fingerprinting (level 2), 237–245
 - biomarkers, 239–240
 - diagnostic ratios, 237–238
 - polycyclic aromatic compounds (PACs), 238–239
 - sesquiterpanes, 240–245
 - sample preparation and cleanup, 233–235
 - treatment of results (level 3), 245–250
 - criteria for diagnostic ratios, 245–246
 - critical difference, 246
 - duplicate analyses, 248–249
 - multivariate statistics, 249–250
 - oil sample comparison, 245
 - repeatability limit, 246
 - signal-to-noise (S/N) test, 246–248
 - visual characterization of samples, 233–235
- CERCLA, 36, 449
- Cerritos Channel, 516
- Cetane rating, 351
- Chain-of-custody documentation, 70
- Charcoal, 39–41
- Charged-coupled devices (CCDs), 421, 439
- Cheilanthane, 242
- Chemical fingerprinting, 257–258
 - control factors, 11–43
 - background chemicals (tertiary controls), 36–43
 - crude oil genesis (primary controls), 13–18
 - petroleum refining (secondary controls), 18–26
 - weathering (tertiary controls), 29–37
 - of heavy fuel oils (HFO), 330–332
 - of known-source spills, 3

- Chemical fingerprinting (*Continued*)
- methods, 3–11
 - GC/FID, 6–7
 - GC/MS, 7–10
 - petroleum biomarkers, 9–10
 - polycyclic aromatic hydrocarbons (PAHs), 8–9
 - historical perspective, 3–6
 - quality assurance, 10–11
 - quality control, 10
 - of mystery spills, 2–3
 - overview, 2–3
 - qualitative, 258–263
 - quantitative, 263–276
- Chemical toxicity, 59–60
- Chevron Biomarker Laboratory, 86
- Chiral carbons, 80–81
- Cholanes, 267
- Cholestane, 14
 - aerobic biodegradation of, 353
 - diagnostic ratios, 243
 - GC/FID analysis of, 267
 - nomenclature, 81
- Chromatograms, 174–175
- ChromatOF (software), 173
- Chrysenes, 5
 - biodegradation of, 361
 - diagnostic indices, 271
 - diagnostic ratios, 237–239
 - GC/FID analysis of, 267
 - in heavy fuel oils (HFO), 339
 - mass fragments, 300
 - in oil spills, 514
 - response factor, 8
- Chrysenothiophene, 149
- Citrobacter freundii*, 357
- Climax communities, 366
- Co-metabolism, 364
- Coal, 39–41
- Coal Oil Point, 196, 211
- Coastal facility spills, 2
- Coastal oil spills, 62–63
- Coelution, 210
- Coenzyme A, 352, 355
- Coenzyme M, 352
- COLD zone, 59
- Columns, 82
- Commercial ships, 328–329
- Commet assay, 390
- Compact airborne spectrographic scanner (CASI), 421
- Compound-specific isotope analysis (CSIA), 208–209, 214–220
- Comprehensive two-dimensional gas chromatography (GC × GC), 171–172
- chromatograms, 174–175
 - data processing, 173–174
 - detectors, 172–173
 - modulation techniques, 172
 - in oil spill fingerprinting, 181–201
 - Buzzards Bay No. 6 spill, 191–195
 - Mobile Bay marine diesel fuel spill, 181–184
 - Santa Barbara oil seep, 195–201
 - West Falmouth fuel spill, 184–187
 - Winsor Cove fuel oil spill, 187–191
 - in oil spill source identification, 158
 - peak identity, 175–180
 - petroleum applications, 180–181
- Compressive waves, 437
- Condensates, 350
- Constitutional isomers, 79
- Constrained least squares (CLS), 466–468
- Continuous-flow instruments, 210
- Conventional crude oils, 13
- Conversion, 20–21
- Correlation optimized warping (COW), 304–305
- Crankcase oil, 494–495
- Critical difference, 246
- Cross-plots, 128–130
- Crude oils, 13–18
 - biodegradation of, 354
 - biomarkers in, 93–99
 - classification of, 13
 - composition of, 13–15, 350
 - density of, 350
 - gas chromatograms of, 359
 - genetically-similar, 260–261
 - vs. heavy fuel oils (HFO), 343–346
 - qualitatively-similar, 261
 - refining, 18–26
 - distillate fuels, 22–27
 - gasoline, 21–22
 - lubricating oils, 28
 - residual fuels, 27–28
 - source markers for, 160–162
 - sulfur compounds in, 148–150
 - ternary diagram for, 13
 - weathering, 29–37
- Cryogenic trapping, 172
- Currents, 411–412
- Cyanobacteria, 365
- Cyclic biomarkers, 81
- Cyclic terpenoids, 75–77
- Cyclododecanol, 352
- Cyclohexane-monoxygenase, 351

- Cyclohexanol, 352
 Cytochrome P₄₅₀, 351, 388–390
- D**
- Dalco Passage oil spill, 63
 Data management, 67–71
 Daughter ions, 85–86
 Daya Bay, 516
 Decahydronaphthalenes, 109, 187–191
 Decalins, 109, 187–191, 267
 Decision chart, 233
 Deep Bay, 516
 Deep subsurface ecology, 366–367
 Delphi 4.0 (software), 296
 Denitrification, 366
 Density, 350
 Density-driven current, 411–412
 Desmethylsteranes, 198–200
 Detectors, 172–173
 Deuterium, 208
 Diacholestane, 243, 267
 Diagnostic indices, 269–272
 Diagnostic power, 126, 313
 Diagnostic ratios, 126–128
 biomarkers, 239–240
 comparison of, 249–254
 criteria for, 245–246, 251–254
 definition of, 232
 elimination of, 246–249
 duplicate analyses, 248–249
 signal-to-noise (S/N) test, 246–248
 from GC/MS fingerprinting, 237–238
 in oil hydrocarbon fingerprinting, 306–307
 oil sample comparison, 245
 polycyclic aromatic compounds (PACs),
 238–239
 polycyclic aromatic hydrocarbons (PAHs),
 510–511
 sesquiterpanes, 240–245
 signal-to-noise (S/N) test, 246–248
 Diahopane, 122–124, 242, 528–530
 Dialkylthiacyclopentanes, 149
 Diamantanes, 78, 114–117, 127
 Diamondoids, 78, 114–117
 Diaromatics, 14
 Diasteranes, 127, 300, 361, 362
 Diastereomers, 79–80
 Dibenzanthracene, 5, 8, 267, 339, 514
 Dibenzofuran, 8, 267, 339
 Dibenzofuranes, 300
 Dibenzothiophenes, 149–150
 biodegradation of, 355, 361
 in crude oils, 152, 161–162
 diagnostic ratios, 237–239
 in distillate fuels, 27
 in *Exxon Valdez* oil spill, 461
 GC/FID analysis of, 267
 in heavy fuel oils (HFO), 339
 mass fragments, 300
 in oil spills, 514
 response factor, 8
 source indices, 271
 Dicarboxylic acid, 352
 Dichloromethane, 266
 Dielectric constants, 437
 Diesel fuel, 7, 22, 99–101, 152, 351
 Diesel range organics (DRO), 38
 Diethylbenzene, 353–355
 Differential global positioning system (DGPS),
 434
 Dihydroxybenzoates, 352
 Dimethylcyclohexane, 356
 Dimethylcyclopentane, 356
 Dimethyldibenzothiophenes, 152
 Dimethylnaphthalenes, 361
 Dimethylphenanthrenes, 361
 Dinaphthothiophene, 149
 Dinosterane, 123
 Dioxin, 394
 Dioxygenases, 351–353
 Diphenyl disulfides, 155
 Discrete biomarkers, 10
 Dispersion, 415
 Dissimilar match, 260
 Dissolution, 32, 131, 382
 Distillate fuels, 22–27
 Distillation, 24, 350–351
 Distribution coefficient, 384
 Diterpanes, 14
 Dodecahydrotriphenylene, 198
 Doppler shifts, 433
 Drimane, 109–111, 244
 Duplicate analyses, 248–249
 Duque de Caxias refinery, 506
 Dynamic time warping (DTW), 304
- E**
- Eagle Harbor, 516
 EC Oil Spill Research Laboratory, 85
 Egyptian crude oil, 161
 Emulsification, 32, 130, 415
 Emulsions, 34
 Enantiomers, 79
 Environment Canada Oil Research Laboratory,
 82–83
 Environmental data, collection of, 63

- ENVISAT satellite, 436
 Enzyme-linked immunosorbent assay (ELISA), 390
 EPA method 8015B, 5
 Ergostanes, 81
Erika oil spill, 33, 217–218, 408
 ERS-1 satellite, 436
 ERS-2 satellite, 436
 Ethane, 361
 Ethene, 352
 Ethoxyresorufin, 390
 Ethyladamantanes, 114–117
 Ethylbenzene, 5
 Ethylbenzene dehydrogenase, 356
 Ethylcholestane derivative, 243
 Ethylmethylbiphenyls, 361
 Ethylphenanthrenes, 361
 Eudesmane, 14, 109
 Eulerian approach, 408
 Evaporation, 30–32, 415
 Excimer laser, 424–425
 Excitation-emission matrices (EEMs), 298, 307–308
 Exposure limits, 59–60
 Extracted ion plots (EIPs), 261
 Extraction, 496
 Extradiol dioxygenases, 352–353
Exxon Valdez oil spill, 449–450. *See also* Oil spills
 - biodegradation in, 153
 - biotic responses to, 396–398
 - composition and chemistry of crude oil in, 453–457
 - forensic investigations, 449–482
 - hydrocarbon source allocation, 462–469
 - qualitative allocation models, 464
 - quantitative models, 464–469
 - source allocation models, 462
 - identification of hydrocarbon sources in, 450–453
 - biogenic hydrocarbons, 453
 - multiple sources, 450–451
 - petrogenic hydrocarbons, 451–453
 - pyrogenic hydrocarbons, 453
 - investigation of hydrocarbons, 470–477
 - oil bioavailability assessment, 477–482
 - biomarkers, 481–483
 - passive sampling of PAH, 481
 - uptake of PAH in biota, 477–480
 - PAH source allocation, 469–470
 - PASHs as markers in, 162–164
 - resolution of natural background, 458–462
 - isotope ratios in, 211–212
 - photooxidation in, 153
 - phytane in, 387
 - polycyclic aromatic hydrocarbons (PAHs) in, 387, 457–458
 - presence of other hydrocarbons in, 13
 - triplet ratios in, 128
 - weathering products, 453–457
 - weathering trends, 457–458
 - data sources, 457
 - major fractions, 457
 - mass loss, 458
 - polycyclic aromatic hydrocarbons (PAHs), 457–458
- F**
- Factor analysis, 294
 Fay, James, 413
 Fermentative bacteria, 369
 Finishing, 20–21
 Flame ionization detectors (FID), 6–7
 Flame photometric detection (FPD), 156–157
 Flash point, 329
 Flasher bottoms, 328
Florida oil spill, 184–187, 449
 Fluoranthenes, 5
 - GC/FID analysis of, 267
 - in heavy fuel oils (HFO), 339
 - mass fragments, 300
 - in oil spills, 514
 - response factor, 8
- Fluorenes, 5
 - GC/FID analysis of, 267
 - in heavy fuel oils (HFO), 339
 - mass fragments, 300
 - in oil spills, 514
 - response factor, 8
- Fluorescence spectroscopy, 298–299, 307–308
 Fluorescent aromatic contaminants (FACs), 481
 Fluvio-deltaic oils, 214
 Forecasting, 406–407
 Forensic investigations, 55–71
 - data collection in, 63–64
 - data management, 67–71
 - emergency response phase, 57–61
 - chemical toxicity analysis, 59–60
 - personal protective equipment (PPE), 60–61
 - risk assessment and characterization, 58–59
 - safety management, 57–58
 - working environment safety, 60
 - geographic boundary determination in, 61–63
 - coastal oil spills, 62–63
 - marine oil spills, 62–63
 - terrestrial spills, 61–62

- identification of hydrocarbon sources in,
 - 450–453
 - biogenic hydrocarbons, 453
 - multiple sources, 450–451
 - pyrogenic hydrocarbons, 453
 - petroleum PASH markers in, 159–164
 - reconnaissance survey in, 55–57
 - sampling plan and design, 64–67
 - known-source spills, 64–66
 - mystery spills, 66–67
 - site characterization in, 55–57
- Fortner Software LLC, 173
- Fractionation, 20–21, 496
- Fraunhofer light discriminator, 424
- Freighter spills, 2
- Friedelene, 90
- Fuel No. 5, 100–101
- Fuel oil, 22, 338–343
- Full scan mode, 84
- Fumarate, 355
- Fungi, 350
- G**
- Gammacerane, 78, 120–121, 267, 528–530
- Gas chromatography/mass spectrometry (GC/MS),
 - 83–86
 - baselines, 303–304
 - benchtop quadropole, 84–85
 - biomarkers, 9–10
 - case study, 251
 - in chemical fingerprinting, 7–10
 - normalization, 306
 - in oil hydrocarbon fingerprinting, 299–301
 - overview, 4–6
 - polycyclic aromatic hydrocarbons (PAHs),
 - 8–9
 - in quantitative fingerprinting, 268–269
 - retention time, 304–305
 - scan mode, 84
 - selected ion monitoring (SIM), 84–85
 - triple quadropole, 85–86
- Gas oils, 22
- Gases, 22
- Gasoline, 21–22, 99, 351
- GC/FID fingerprinting, 6–7
 - case study, 251
 - in CEN oil spill identification, 235–237
 - in quantitative fingerprinting, 266–268
- GC Image (software), 173
- Gelbstoff, 424
- Gene regulation, 364
- Geobacter*, 368
- Geographic information system (GIS), 61, 68
- Ghawar oil field, 357
- Gironde Bay, 516
- Global positioning system (GPS), 68, 420
- Gram-negative bacteria, 356
- Gravity, 350
- Groundwater, 161
- Guanara Bay, 506
- Guanara Bay oil spill, 506–507
 - chemical analysis, 508–510
 - biomarkers, 509–510
 - extract cleanup, 509
 - PAH analysis, 508–510
 - sediment sample extraction, 508–509
 - source identification methods, 508–510
 - hydrocarbon source identification, 514–532
 - biomarker diagnostic ratios, 527–532
 - PAH diagnostic ratios, 514–525
 - PAH principal component analysis,
 - 525–527
 - PAH quantification, 511–514
 - sampling design, 507–508
 - source identification methods, 510–511
 - biomarker diagnostic ratios, 511
 - PAH diagnostic ratios, 510–511
 - PAH multivariate analysis, 511
- Gulf of Suez crude oil, 161
- Gullfaks oilfield, 367
- H**
- Halophilic bacteria, 364
- Haven* oil spill, 162, 435
- Heavy crude oils, 13
- Heavy fuel oils (HFO), 327–346. *See also* Crude oils; Petroleum
 - boiling points, 332–333
 - chemical fingerprinting, 330–332
 - classification of, 329–330
 - composition of, 333–336, 351
 - vs. crude oil, 343–346
 - historical perspective, 328
 - ISO specifications for, 329
 - isoalkane ratios, 331
 - molecular variability in, 336–343
 - biomarkers, 336–338
 - polycyclic aromatic hydrocarbons (PAHs),
 - 338–343
 - overview, 327
 - production of, 329–330
 - quantitative fingerprinting of, 279–289
 - samples and analytical methods, 332
 - total hydrocarbons, 331
- Heavy residual fuels, 101
- Heavy straight-run distillates, 22

- Heliscat, 428
 Hematite, 368
 Heptadecane, 236, 352
 Herring, 474–475
 Heterocyclic aromatic compounds, 300
 Hexadecane, 352, 357
 Hindcasting, 406–407
 Holly oil, 196–201
 Homodrimanes, 110, 244
 Homohopanes, 201, 242, 267
 Homopregnane derivative, 243
 Hopanes, 80
 analysis of, 496–497
 biodegradation of, 133
 aerobic, 353
 subsurface, 360–361
 susceptibility, 362
 as biomarkers, 124, 199–201
 diagnostic indices, 271
 diagnostic ratios of, 242, 528–530
 GC/FID analysis of, 267
 as internal standards, 134
 isomers, 81
 mass spectra of, 87
 structures of, 14
 HOT zone, 59
 Houston Ship Channel, 211
 Huntington Beach, 212
 Hurricane Katrina, 31, 38–39
 Hydrocarbons, 349–350
 accumulation by biota, 382–387
 aerobic biodegradation of, 351–355
 aliphatic, 14
 anaerobic biodegradation of, 355–357
 aromatic, 14
 biogenic, 350, 451, 453
 branching, 352
 catabolic degradation of, 387–393
 identification of, 381–382, 470–477
 labeling and nomenclature of, 77–81
 petrogenic, 350, 450–453
 pyrogenic, 451, 453
 structures of, 74–77
 cyclic terpenoids, 74–77
 isoprenoids, 74
 toxicity of, 393–396
 Hydrodesulfurization, 27, 151
 Hydrodynamic models, 412
 Hydrogen isotopes, 208, 224
 Hydrothermal vents, 365
 HyperChrom (software), 173
 Hyperspectral imaging, 421
 Hyperthermophilic organism, 365, 368
- I**
 Ice, 436–438
 Inconclusive match, 260
 Indan, 177
 Indene, 177
 Indenopyrene, 5, 8, 267, 339
 Indeterminate match, 259
 Industrial Materials Institute (NMI), 433
 Infrared cameras, 438–439
 Infrared sensors, 422–423
 Integrated multivariate oil fingerprinting (IMOF), 296–297
 Intentional operational discharges, 1–2
 Intermediate fuel oil 180 (IFO 180), 100–101, 113, 330
 Intermediate fuel oil 380 (IFO 380), 330
 biomarkers, 336–338
 chromatograms, 333–336
 polycyclic aromatic compounds (PACs), 338–342
 Intermediate fuel oil (IFO), 279–289
 Internal injection standard (IISTD), 497
 Internal standards, 83
 Intradiol dioxygenases, 352–353
 Iron-reducing bacteria, 368
 Iron reduction, 366
Iso-butane, 361
Iso-paraffins, 14
 ISO specifications, 329–330
 Isoalkane ratios, 331
 Isoalkanes, 361
 Isoprenes, 74
 Isoprenoids, 7, 75
 biodegradation of, 361
 diagnostic indices, 271
 diagnostic ratios, 127, 253
 mass fragments, 300
 susceptibility to biodegradation, 362
 Isopropyl benzene, 354–355
 Isotope ratio mass spectrometer (IRMS), 208–210
 Isotopes, 208
 Isowake Experiments, 431
- J**
Jacob Luckenbach, 67
 Jet fuel, 99
 Jiulongjiang Estuary, 516
 Johnston Creek crude oil, 460–461
- K**
 Katalla Formation shale, 459–462
 Kendrick mass defect, 154
 Kendrick nominal mass, 154

- Kerogens, 349, 467
 Kerosene, 22, 113
 Ketones, 352
 Known-source spills, 3, 64–66
 Knudsen Engineering Ltd., 437
 Korea Diesel No. 2, 100–101, 110
 Kulthieth Formation, 456–457
Kurdistan oil spill, 435
 Kuskutz coals, 456, 459–461
 Kuwait crude oil, 90
- L**
- L-band radar, 427
 La Luna crude oil, 359
 Lactone, 352
 LANDSAT satellite, 435
 Langragian elements, 408
 Laser fluorosensors, 419–420, 423–426, 438–439
 Laser ultrasonic remote sensing of oil thickness (LURSOT), 433–435
 Leco Corp., 173
 Lewis acids, 114
 Light distillates, 99
 Light oils, 350
 Light straight-run distillates, 22
 Lignite, 40
 Lipid catabolic pathways, 352
 Loading plots, 315–317
 Low-density polyethylene strips (LDPEs), 481
 Lowest effect level (LEL), 59
 Lubricating oils, 28
 - biomarkers, 101–103
 - boiling point, 22
 - carbon range, 22
 - use in marine fuels, 328–329
- M**
- Macrocyclic alkanes, 124–125
 Malaysia, oil pollution sources in, 491–492
 Manganese reduction, 366
 Manmade petroleum, 12
 Marine carbonate oils, 214
 Marine diesel oil (MDO), 279, 284–286, 330
 Marine Ecosystems Research Laboratory (MERL), 187
 Marine fuel oils, 327–346. *See also* Crude oils; Petroleum
 - boiling points, 332–333
 - chemical fingerprinting, 330–332
 - classification of, 329–330
 - composition of, 333–336
 - vs. crude oil, 343–346
 - historical perspective, 328
 - ISO specifications for, 329–330
 - isoalkane ratios, 331
 - molecular variability in, 336–343
 - biomarkers, 336–338
 - polycyclic aromatic hydrocarbons (PAHs), 338–343
 - overview, 327
 - production of, 329–330
 - quantitative fingerprinting of, 279–289
 - samples and analytical methods, 332
 - total hydrocarbons, 331
- Marine gas oil (MGO), 330
 Marine oil spills, 57, 62–63, 181–184, 277–279, 327
 Masan Bay, 516
 Mass fragments, 300
 Mass losses, 458
 Mass-selective detection (MSD), 157
 Mass selective detector (MSD), 85
 Mass spectrometry, 4–6
 Mass-to-charge ratio, 84–90
 Match, 259
 Math Works Inc., 173
 Matlab (software), 173, 296
 Matrices, 310–311
 Maximum constructive interferences, 429
 Maximum destructive interferences, 429
 Mercaptans, 350
 Mersey Estuary, 516
 MESA oil, 105–107
 Mesozoic petroleum system, 17
 Methane, 361
 Methanogenesis, 357, 366–367
 Methanogens, 356–357, 369
 Method detection limits (MDLs), 465–466
 Methyldiamantane index (MDI), 116
 Methyl-ethyl benzene, 354–355
 Methyl groups, 90, 155
 Methyldadamantane index (MAI), 116
 Methyldadamantanes, 114–117
 Methylanthracene, 267
 Methylation, 355
 Methylbiphenyls, 361
 Methylcholestanes, 243, 267
 Methylchrysene, 267
 Methyl dibenzothiophenes, 152, 237–239, 267, 296
 Methylfluoranthenes, 238
 Methylfluorene, 295
 Methyl naphthalenes, 222, 361, 514
 Methylpentylsuccinate synthase, 355
 Methylphenanthrenes, 237–239, 267, 271, 296, 361

- Methylpyrenes, 237–239
Methylsteranes, 124
Metula oil spill, 133, 450
Microaerophilic bacteria, 368
Microbial communities, succession of, 365–367
Microbial ecology, 365
Microcarbon residues, 329
Microwave sensors, 426–428. *See also* Remote sensing
 radar, 426–428
 radiometers, 426
 scatterometers, 428
Mid-range distillates, 99–101
Midcontinent rift system, 367
Minas oil, 216
Minjiang Estuary, 516
Miocene Monterey Formation, 451
Mirror-image isomers, 81
Mixtures, separation of, 169–170
Mobile Bay marine diesel fuel spill, 181–184
Models and modeling, 405–416
 currents, 411–412
 oil spill movement, 406–407
 oil spill transport, 407–409
 overview, 405–406
 spreading, 413–414
 turbulence, 413
 weathering, 414–416
 wind, 409–411
Modulators, 172
Molecular fossils. *See* Biomarkers
Molecular markers, 501–502
Molybdopterin cofactor, 356
Monoaromatic steranes, 127
Monoaromatics, 14
Monocyclic acids, 361
Monomethylbenzothiophenes, 155
Monooxygenases, 351, 354
Monterey Formation, 128
Moretane, 242, 267, 271
Mousse, 34
Multi-detector electrooptical imaging scanner (MEIS), 421
Multi-linear models, 310–312
 higher-order arrays, 311–312
 two-way case, 310–311
Multi-spectral scanners, 438–439
Multidimensional separation, 170–171
Multivariate statistics, 308–314
 in comparison of diagnostic ratios, 249–250
 constrained least squares (CLS), 466–468
 multi-linear models, 310–312
 in oil spill fingerprinting, 294–296
 partial least squares (PLS), 468
 variable selection and scaling, 312–314
Mussels, 387, 391–393, 477–480
Mycobacterium vanbaalenii, 351
Mysidium gracile, 507
Mystery spills, 2–3, 66–67
- N**
N-alkanes, 90, 361, 362, 383
N-butane, 361
N-cyclohexyl-2-benzothiozalamine (NCBA), 497
N-way toolbox, 297
Naphthalene dioxygenase, 352
Naphthalenes, 339
Naphthenes, 14, 350
Naphthenic acids, 361
Naphthobenzothiophenes, 8, 267, 271, 300
Naphthothiophenes, 27, 149
Naphthalene carboxylase, 356
Naphthalenes, 5
 carbon isotope ratios, 222
 GC/FID analysis of, 267
 mass fragments, 300
 in oil spills, 8, 183, 514
 vapor pressure of, 382
National Petroleum Reserve, Alaska (NPRA), 367
Natural dispersion, 130–131
Natural hydrocarbons, 37–41
 algal debris, 38–39
 oil seeps, 41
 vascular plants, 38–39
 wood charcoal, 39–41
Natural resource damage assessment (NRDA), 265
Navy Fuel Oil (NFO), 67
Nd:YAG laser, 424, 433
NetCDF (software), 296
New Carisa, 68
Nigerian Bonnie Light crude oil, 16–17
Night vision cameras, 423
Nipisi oil spill, 90
NIST 1582 oil, 110
Nitrate-reducing bacteria, 366
Nitrogen isotope ratios, 211, 224
Nitrogen laser, 424
Nitrogen oxides, 150
Nomenclature, 77–81
Non-tanker spills, 2
Nonane, 353, 383
Nonaqueous phase liquids (NAPLs), 99
Nonmatch, 251, 259, 273, 319
Nordtest method, 229–230, 273–276
Norhomohopane, 528–530

- Norhopanes, 360–361
 biomarker indices, 271
 diagnostic ratios, 242
 formation of, 360–361
 GC/FID analysis of, 267
 mass spectra of, 87
 susceptibility to biodegradation, 362
- Normalization, 306
- Normoretane, 242, 267
- Norneohopane, 133, 242, 267
- Norpristane, 267, 271
- O**
- Ocean circulation, 411–412
- Octadecane, 236, 352
- Octane monooxygenase, 351
- Octane rating, 351
- Office of Solid Waste and Emergency Response, 36
- Offshore rigs, discharges from, 2
- Oil embargos, 328
- Oil fractions, 104, 213, 457
- Oil hydrocarbon fingerprinting, 293–294. *See also*
 Biomarker fingerprinting
 analytical methods, 298–300
 fluorescence spectroscopy, 298–299
 gas chromatography/mass spectrometry (GC/MS), 299–301
 data evaluation, 314–319
 numerical comparisons and statistical tests, 317–319
 score and loading plots, 315–317
 data preprocessing, 302–308
 diagnostic ratios, 306–307
 fluorescence spectra, 307–308
 partial GC-MS/SIM chromatograms, 303–306
 baseline removal, 303–304
 normalization, 306
 retention time alignment, 304–305
 integrated multivariate, 296–297
 multivariate statistics, 294–296, 308–314
 multi-linear models, 310–312
 higher-order arrays, 311–312
 two-way case, 310–311
 variable selection and scaling, 312–314
 quality assurance, 301–302
 quality control, 301–302
 sample preparation, 297
- Oil-in-ice detectors, 436–438
- Oil-mineral aggregation, 132
- Oil production platform spills, 2
- Oil seeps, 1, 41, 196–201
- Oil spills, 1–3
 accidental, 1–2
 chemical toxicity of, 59–60
 coastal, 62–63
 evolution of, 413–416
 oil weathering, 414–416
 spreading, 413–414
 fingerprinting of, 181–201
 Buzzards Bay No. 6 fuel spill, 191–195
 Mobile Bay marine diesel fuel spill, 181–184
 Santa Barbara oil seep, 196–201
 West Falmouth No.2 fuel oil spill, 184–187
 Winsor Cove fuel oil spill, 187–191
 identification of, 231–233
 intentional operational discharges, 1–2
 known-source release, 3
 on land, 56
 marine, 62–63
 modeling of, 405–416
 forecasting and hindcasting, 406–407
 overview, 405–406
 transport, 407–413
 currents, 411–412
 turbulent diffusion, 413
 wind, 409–411
 multidimensional analysis of, 170–171
 mystery spills, 2–3
 remote sensing of, 419–439
 acoustic systems, 435
 detection of oil under ice, 436–438
 future trends in, 438–439
 laser fluorosensors, 423–426
 microwave sensors, 426–428
 radar, 426–428
 radiometers, 426
 scatterometers, 428
 optical sensors, 420–423
 infrared, 422–423
 night vision cameras, 423
 ultraviolet, 423
 visible, 420–422
 visible indications of oil, 420–422
 real-time displays and printers, 438
 satellite, 435–436
 slick thickness, 428–435
 remote sensors, 431–432
 specific sensors, 431–435
 visual indicators, 428–431
 visible indications of oil, 420
 sensitivity of birds to, 216–217
 source identification of, 125–130, 134–138
 sources of, 1–2

- Oil spills (*Continued*)
 terrestrial, 61–62
 on water, 56
- Oil-water interface, 429
- Oily wastes, 28
- Oleanane, 90
 diagnostic indices, 271
 diagnostic ratios, 242, 528–530
 GC/FID analysis of, 267
 identification of, 459
 in oil spills, 120–123
 structure of, 14
- Operational discharges, 1–2
- Optech Inc., 426
- Optical frequency compensation device (OFCD), 433
- Optical sensors, 420–423. *See also* Remote sensing
 infrared, 422–423
 night vision cameras, 423
 ultraviolet, 423
 visible, 420–422
- Organo-sulfur compounds, 148–150
- Orimulsion, 425
- Orinoco crude oil, 96–99, 357
- Oscillatoria*, 365
- Oserberg Field Centre oil, 315–317
- Oserberg South East oil, 315–317
- Oxidants, 365–366
- Oxidation, 415–416
- Oxidative element analyzers, 210
- Oxidative metabolic gearing, 368
- Oxygen, 351–352
- Oxygen isotope ratios, 224
- P**
- Paraffins, 14, 350, 359
- Parallel factor analysis (PARAFAC), 307–312
- Parent ions, 85–86
- Partial least-squares regression, 295, 310, 468
- Particles, 408–409
- Particulate coal, 39–41
- Pattern recognition, 294
- Pearl River Estuary, 516
- Pee Dee Belemnite (PBD), 208
- Pentaaromatics, 14
- Pentacyclic terpanes, 135–138
- Pentacyclic triterpanes, 267
- Pentakishomohopanes, 267
- Pentanes, 361
- PERF Method, 62
- Perfluorotributylamine (PFTBA), 85, 268
- Personal protective equipment (PPE), 60–61
- Perylene, 40
 GC/FID analysis of, 267
 generation of, 453
 in heavy fuel oils (HFO), 339
 in oil spills, 8, 514
- Petrobras, 506
- Petrobras Geochemistry Laboratory, 85
- Petrogenic hydrocarbons, 350, 450–453
- Petroleum, 11–12. *See also* Crude oils
 aromatic steranes in, 104–109
 biomarkers in, 99–101
 chemical fingerprinting of, 258
 crude oil genesis (primary control), 13–18
 high-resolution separation of, 169
 manmade, 12
 natural, 11
 primary controls, 12
 refining, 18–26
 distillate fuels, 22–27
 gasoline, 21–22
 lubricating oils, 28
 residual fuels, 27–28
 secondary controls, 18–26
 sulfur compounds in, 148–150
 tertiary controls, 29–37
 weathering, 12, 29–37
- Petroleum biodegradation. *See* Biodegradation
- Petroleum biomarkers. *See* Biomarkers
- Petroleum products, 350–351
- Phenanthrenes, 5
 biodegradation of, 355, 361
 diagnostic ratios, 237–239
 in distillate fuels, 27
 in *Exxon Valdez* oil spill, 461
 in heavy fuel oils (HFO), 339
 mass fragments, 300
 in Mobile Bay marine fuel spill, 267
 in oil spills, 183–184, 514
 response factor, 8
- Phenanthrothiophene, 149
- Phenazines, 369
- Phenols, 360, 361
- Phosphorus, 329
- Photochemical processes, 155
- Photooxidation, 34, 131–132, 153, 382
- Physical data, collection of, 63
- Phytane, 75, 236, 267, 352
- Pink salmon, 474–477
- Pipeline spills, 2
- Plant biomarkers, 90
- Platform Elly crude oil, 96–99, 109
- Point sources, 1
- Point-to-point matching, 295

- Polars, 15, 360
- Polycyclic aromatic compounds (PACs), 160, 232–233, 238–239, 300
- Polycyclic aromatic hydrocarbons (PAHs), 8–9.
See also Biomarkers
 accumulation of, 390–393
 aerobic biodegradation of, 355
 analysis of, 497–498
 catabolic degradation of, 387–393
 depuration of, 390–393
 diagnostic indices, 269–272
 diagnostic ratios, 510–511, 514
 distribution coefficient of, 384
 in *Exxon Valdez* oil spill, 457–458
 in heavy fuel oils (HFO), 338–343
 in light oils and condensates, 350
 mass fragments, 300
 multivariate analysis of, 511
 persistence of, 390–393
 principal component analysis (PCA), 514
 pyrogenic index, 457–458
 in qualitative fingerprinting, 264–265
 quantitative evaluation of, 135–138
 in quantitative fingerprinting, 268, 464–465
 separation from PASHs, 157–158
 source allocation of, 469–470
 susceptibility to biodegradation, 362
 uptake in biota, 477–480
 in urban runoff, 42–43
- Polycyclic aromatic sulfur heterocycles (PASHs), 147–164. *See also* Biomarkers
 analysis methods for, 155–159
 selective detection in gas chromatography, 156–157
 atomic emission detection (AED), 157
 flame photometric detection (FPD), 156–157
 mass-selective detection (MSD), 157
 sulfur chemiluminescence detection (SCD), 157
 separation from PAH, 157–158
 two-dimensional gas chromatography (GC x GC), 158
 in Egyptian crude oil, 152
 identification of, 150
 influence of refinery processes on, 150–152
 overview, 147–148
 quantification of, 158–159
 as source markers, 160–162
 stability of, 152–155
 structures of, 148–150
 as weathering markers, 162–163
- Polydimethylsiloxane, 174
- Polysilphenylene, 174
- Porphyrins, 125, 360, 361, 362
- Portland Harbor, 516
- Positive match, 250–251, 273, 319
- Potentially responsible party (PRP), 230, 265
- Poul Creek crude oil, 460–461
- Pour point, 329
- Precambrian period, 350
- Pregnane derivative, 243
- Prestige* oil spill, 428
- Primary controls, 13–18
- Prince William Sound, 41, 128, 211–212, 295, 450–453
- Principal component analysis (PCA), 249–250, 294–296, 310, 465–466
- Principal component regression (PCR), 310
- Printers, 438
- Priority pollutants, 5
- Pristane, 14
 biodegradation of, 352
 biomarker indices, 271
 diagnostic ratios, 236
 GC/FID analysis of, 267
- Probable match, 251, 259, 273, 319
- Procedural artifacts (PA), 471–474
- Propane, 361
- Propyl benzene, 354–355
- Propylene, 352
- Proteobacteria, 356–357
- Prudhoe Bay oil, 115
- Pseudomonads, 364
- Pseudomonas*, 153
- Pseudomonas mendocina*, 354
- Pseudomonas putida*, 351–354
- Pseudomonas stutzeri*, 353
- Pyrenes, 5
 GC/FID analysis of, 267
 in heavy fuel oils (HFO), 339
 mass fragments, 300
 in oil spills, 514
 response factor, 8
- Pyridines, 361
- Pyrite, 364
- Pyrogenic hydrocarbons, 451, 453
- Pyrogenic index, 457–458
- Pyrrols, 361
- Q**
- Qualitative allocation models, 464
- Qualitative fingerprinting, 258–263
 match criteria, 259–260
 shortcomings of, 260–263
 genetically similar oils, 260–261

- Qualitative fingerprinting (*Continued*)
 mixing, 261–263
 qualitatively similar oils, 261
 weathered oils, 260
- Quality Assurance Project Plan (QAPP), 70
- Quality assurance (QA), 10–11, 70–71, 301–302
- Quality control (QC), 10, 70–71, 301–302
- Quality management plan (QMP), 10
- Quantitative fingerprinting, 263–265
 in analysis of mixed source oil, 276–289
 fuel oil spills, 279–289
 marine oil spills, 277–279
 two-component mixing models, 276–277
- data generation for, 265–269
 data quality, 269
 GC/FID analysis, 266–268
 GC/MS analysis, 268–269
 sample collection, 265–266
 sample preparation, 266
- diagnostic indices, 269–272
 source identification protocols, 272–276
- Quantitative models, 464–469
 PAH ratios, 464–465
 statistical models, 465–468
 constrained least squares (CLS), 466–468
 partial least squares (PLS), 468
 total organic carbon (TOC), 468–469
- Queen Vein coals, 460–461
- Quinones, 369
- R**
- Radar, 420, 426–428, 438
- Radar satellites, 436
- Radarsat satellite, 436
- Radiolysis, 352
- Radiometers, 426, 438–439
- Ralstonia pickettii*, 351
- Raman scattering, 298–299, 424
- Rayleigh scattering, 298–299
- Real-time displays, 438
- Recent organic matter (ROM), 469
- Recreational marine vessel, discharges from,
 2
- Reductive element analyzers, 210
- Reference index (RI), 114
- Refineries, 20–21, 350–351
- Refractive index, 429, 459
- Relative response factors (RRFs), 9, 266
- Relative standard deviation (RSD), 249, 313
- Remote sensing, 419–439
 acoustic systems, 435
 future trends in, 438–439
 microwave sensors, 426–428
 radar, 426–428
 radiometers, 426
 scatterometers, 428
 optical sensors, 420–423
 infrared, 422–423
 night vision cameras, 423
 ultraviolet, 423
 overview, 419–420
 real-time displays and printers, 438
 satellite, 435–436
 slick thickness, 428–435
 remote sensors, 431–432
 specific sensors, 431–435
 visual indicators, 428–431
 visible indications of oil, 420
- Remotec Applications Ltd., 436
- Repeatability limit, 246
- Reservoir biodegradation, 357–358
- Reservoir strata, 11
- Residual fuel oils, 327–346. *See also* Crude oils;
 Petroleum
 boiling points, 332–333
 chemical fingerprinting, 330–332
 classification of, 329–330
 composition of, 333–336
 vs. crude oil, 343–346
 historical perspective, 328
 ISO specifications for, 329–330
 isoalkane ratios, 331
 molecular variability in, 336–343
 biomarkers, 336–338
 polycyclic aromatic hydrocarbons (PAHs),
 338–343
 overview, 327
 production of, 329–330
 quantitative fingerprinting of, 279–289
 samples and analytical methods, 332
 total hydrocarbons, 331
- Residual fuels, 27–28
- Residuals, 328
- Resolution power, 312
- Resonance scattering, 437
- Resorufin, 390
- Retene, 237–239, 267, 271, 453
- Retention time, 151, 304–305
- Retention time markers, 10
- % Ring number, 271
- Rio de la Plata Estuary, 516
- Risk assessment, 58–59
- River runoff, 42–43
- Rotational freedom, 310
- Round Robin exercise, 230
- Runoff, 42–43

S

- Safety management, 57–58
- Safety response zones, 58–59
- Sampling, 64–67
 - design, 65–66
 - known-source spills, 64–66
 - on land, 65
 - oil source, 65
 - water column, 64–65
- San Francisco Bay, 516
- Sandstones, 368
- Santa Barbara oil seep, 196–201, 405
- Sarasota Bay, 516
- Satellite remote sensing, 435–436
- Scanners, 421
- Scatterometers, 428
- Scores, 315–317
- Scotia Light crude oil, 93–99, 111–113
- Secondary biogenic methane, 369
- Secondary controls, 18–35
- Sedimentation, 132
- Sediments, 492–494
- Seepage, 195–201
- Select ion monitoring (SIM), 268, 332
- Selected ion monitoring (SIM), 7, 84–85
- Semipermeable membrane devices (SPMDs), 220, 481
- Separation methods, 169–170
- Sequence of priority, 81
- Sesquiterpanes, 109–114. *See also* Biomarkers
 - as biomarkers, 24
 - characterization of, 135
 - cross-plots of, 129–130
 - diagnostic ratios, 127, 137, 240–245
 - labeling and nomenclature of, 78
 - mass fragments, 300
 - structures of, 14
- Seychelles Islands, 218
- Shear waves, 437
- Shipborne radars, 428
- Shoreline Cleanup Assessment Technique (SCAT), 62
- Side-looking airborne radar (SLAR), 427
- Signal-to-noise (S/N) test, 246–248
- Silica gel, 82
- Similar match, 259–260
- Singular value decomposition (SVD), 310
- SINTEF, 231
- Site investigations, 55–71
 - data collection in, 63–64
 - data management, 67–71
 - emergency response phase, 57–61
 - chemical toxicity analysis, 59–60
 - personal protective equipment (PPE), 60–61
 - risk assessment and characterization, 58–59
 - safety management, 57–58
 - working environment safety, 60
- factors in, 57
- geographic boundary determination in, 61–63
 - coastal oil spills, 62–63
 - marine oil spills, 62–63
 - terrestrial spills, 61–62
- petroleum PASH markers in, 159–164
- reconnaissance survey in, 55–57
- sampling plan and design, 64–67
 - known-source spills, 64–66
 - mystery spills, 66–67
- site characterization in, 55–57
- Sockeye crude oil, 96–99
- Sohxlet extraction, 219, 297
- Source allocation models, 462
- Source ratios, 160
- Source-specific biomarkers, 121–125. *See also*
 - Biomarkers
 - 4-methyl steranes, 124
 - beta-carotane, 124
 - bicadinanes, 124
 - diahopane, 122–124
 - hopanes, 124
 - macrocyclic alkanes, 124–125
 - polycyclic aromatic sulfur heterocycles (PASHs), 160–162
 - porphyrins, 125
- Source strata, 11
- South Louisiana crude oil, 93–99
- Southern California Monterey Formation, 212
- Specific gravity, 350
- Sphalerite, 364
- Sphingomonas macrogolitabida*, 353
- Sphingomonas yanoikuyae*, 354
- Spills. *See also* Oil spills
- Spreading, of oil spills, 413–414
- Squalane, 75
- St. Lawrence River, 218
- Stable isotope ratios, 207–224
 - basis of, 208
 - bulk isotope ratios, 210–213
 - compound-specific isotope analysis (CSIA), 214–220
 - in geochemical materials, 209
 - instruments, 208–210
 - measurement of, 207–210
 - overview, 207
 - standards, 208
 - weathering, 220–224
- Standard Mean Ocean Water (SMOW), 208

- Steranes, 75–77. *See also* Biomarkers
biodegradation of, 133, 353, 361–362
diagnostic indices, 271
diagnostic ratios, 127, 137, 528–530
evaluation of, 135–138
in *Exxon Valdez* oil spill, 461
labeling and nomenclature of, 78
mass fragments, 300
mass spectra of, 86–90
structures of, 14
- Stereoisomers, 79–80
- Steroids, 240
- Stigmastanes, 81, 133–134
- Still cameras, 438–439
- Strait of Malacca, 489
hydro-oceanographic conditions of, 489–491
oil pollution sources in, 491–492
oil spill events in, 491
oil spill investigations, 492–502
analytical procedure, 495–496
biomarkers, 498–501
instrumental analysis, 496–498
alkanes, 496–497
hopanes, 496–497
N-cyclohexyl-2-benzothiazolamine
(NCBA), 497
polycyclic aromatic hydrocarbons (PAHs),
497–498
methodology, 492–495
aerosol samples, 495
asphalt samples, 494
extraction, 496
fresh crankcase oil, 494
sample collection, 492
sediment samples, 492–494
source petroleum, 492
street dust samples, 493
tar-ball samples, 492
tire rubber, 495
used crankcase oil, 495
molecular markers, 501–502
polycyclic aromatic hydrocarbons (PAHs),
502
ship traffic in, 491
- Street dust, 493
- Subsurface biodegradation, 357–362
- Subsurface circulation models, 412
- Succinates, 355
- Suez-Mediterranean pipelines, 163
- Sulfate-reducing bacteria, 357, 366–367
- Sulfate reduction, 366
- Sulfides, 148
- Sulfolobus solfataricus*, 365
- Sulfur, 148
- Sulfur chemiluminescence detection (SCD),
157
- Sulfur compounds, 26–27, 148–150, 214
- Sulfur-free gasoline, 150
- Sulfur isotope ratios, 211, 213, 224
- Sulfur oxides, 150
- Surrogates, 83
- Synechococcus*, 365
- Synthetic aperture radar (SAR), 427
- Synthetic microorganisms, 356
- T**
- Tamar Estuary, 516
- Tanker spills, 2
- Tar balls, 69, 120, 211–213, 492, 499–502
- Tar sands, 357
- Tensors, 311
- Tenyo Maru* oil spill, 422
- Teresso industrial oil, 110, 113
- Terminal electron acceptors, 362–363, 366, 369
- Terpanes, 75
biodegradation of, 133
diagnostic ratios, 127, 137, 528–530
evaluation of, 135–138
in *Exxon Valdez* oil spill, 461
labeling and nomenclature of, 78
mass fragments, 300
- Terpenes, 74
- Terpenoids, 74–77
acyclic, 75
cyclic, 75–77
- Terrestrial spills, 61–62
- Tertiary controls, 29–43
background chemicals, 37–43
weathering, 29–36
- Tertiary petroleum system, 17
- Tetraaromatics, 14
- Tetrakishomohopanes, 267
- Tetralin, 353
- Tetramethyladamantanes, 114–117
- Tetramethylphenanthrenes, 361
- TFE-fluorocarbon polymer, 265
- Thaurea aromatica*, 355
- Thermal mapping, 432–433
- Thermo Electron Corp., 173
- Thermopiles, 364–365
- Thermus*, 364
- Thiacyclohexane, 149
- Thiamonocyclanes, 148
- Thiapolycyclanes, 148
- Thiopenes, 149
- Thiopenes, 149–150

Three percent (3%) rule, 410
 Tidal current, 411–412
 Time-of-flight mass spectrometry (TOFMS), 173
 Tire rubber, 495
 Toluene, 5, 14, 353, 364
 Toluene-3-monooxygenase, 351
Torrey Canyon, 405
 Total ion chromatogram (TIC), 84
 Total organic carbon (TOC), 459, 468–469
 Total petroleum hydrocarbons (TPH), 38–39, 266
 Toulon Harbor, 516
 Toxicity, 59–60
 Transform (software), 173
 Transmittance, 424
 Triaromatic steranes, 271, 300
 Triaromatic steroids, 243, 267
 Triaromatics, 14
 Tricyclic terpanes, 361
 Tricyclic triterpanes, 267
 Trimethyl benzene, 354–355
 Trimethylbiphenyls, 361
 Trimethylnaphthalenes, 361
 Triple quadrupole GC-MS-MS, 85–86
 Triplet ratios, 128
 Trishomohopanes, 242, 267
 Trisnorhopane, 242, 267
 Trisnorneohopane, 242
 Triterpanes, 14
 diagnostic indices, 271
 diagnostic ratios, 240–244, 253
 distribution of, 120
 GC/FID analysis of, 267
 Troll crude oil, 93–99
 Turbulence, 413
 Two-component mixing models, 276–277

U

Ultraviolet cameras, 438
 Ultraviolet sensors, 423
 Uncertainty analysis, 409
 Unresolved complex mixture (UCM), 359
 in GC/FID analysis, 7, 42–43
 in marine fuel oils, 333–334
 in oil spills, 184–186
 Urban runoff, 42–43
 Ursene, 90

V

Vacuum gas oils, 22
 Van der Waals force, 382
 Vanadium, 329

Vapor pressure, 382
 Vapors, 59
 Variable-outlier detection, 313
 Variables, 312–314
 Vascular plants, 38–39
 Vegetation biomarkers, 90
 Venezuela Fuel No. 6, 100–101
 Vertical velocity sensor, 434
Vibrio fischeri, 507
 Video cameras, 421, 438–439
 Viscosity, 329, 351
 Visible spectrum, 420–422
 Volatile organic compounds, 60
 Volatile solvents, 22

W

Wallach, Otto, 74
 WARM zone, 59
 Warping path, 304–305
 Water-soluble fraction (WSF), 382, 471
 Water-washing, 384
 Waterborne oil spills, 62–63
 Wavelengths, 429
 Waxes, 34–35
 Waxy bitumens, 215
 Weathered percentages, 134
 Weathering, 29–37
 biodegradation. *See* Biodegradation
 de-waxing, 34–35
 definition of, 12
 dissolution, 32, 131
 emulsification, 130
 evaporation, 30–32, 130
 GC/FID screening, 236–237
 and hydrogen accumulation by biota, 382–383
 markers, 162–163
 and mass loss, 458
 mousse formation, 34
 natural dispersion, 130–131
 oil-mineral aggregation, 132
 in oil spills, 414–416
 photooxidation, 34, 131–132
 and qualitative fingerprinting, 260
 sedimentation, 132
 and stable isotope ratios, 220–224
 wax enrichment, 34–35
 Weathering ratios, 160
 Weighted least-squares analysis, 295, 312–314
 West Falmouth fuel spill, 184–187
 West Texas crude oil, 111–113
 Wind, 409–411
 Wind drift factor, 410
 Wind-driven current, 411–412

Winsor Cove fuel oil spill, 187–191
Working environment, safety of, 60

X

X-band radar, 427–428
XeCl excimer laser, 424–425
Xylene monooxygenase, 354
Xylenes, 5

Y

Yakataga Formation shale, 460–461
Yalujiang Estuary, 516

Z

Zinc, 329