
Index

A

- Acadian flycatcher (*Empidonax virescens*), 604, 606
- Adaptive Resource Management (ARM), models, 11–12
- ALCES, *see* A Landscape Cumulative Effects Simulator
- Allometric scaling relationships, multispecies conservation planning, 53
- ALMaSS, *see* Animal, Landscape, and Man Simulation System
- American bittern (*Botaurus lentiginosus*), 534
- American coot (*Fulica americana*), 534
- American crow (*Corvus brachyrhynchos*), 634, 640–641, 659
- American goldfinch (*Carduelis tristis*), 658
- American robin (*Turdus migratorius*), 658
- American white pelican (*Pelecanus erythrorhynchos*), 534
- American woodcock (*Scolopax minor*), 215, 380
- Animal, Landscape, and Man Simulation System (ALMaSS), habitat-relationships modeling, 257, 265, 269
- Anna's hummingbird (*Calypte anna*), 659
- Arc Habitat Suitability Index (ArcHSD), habitat-relationships modeling, 257, 268
- Arc-Habcap, habitat-relationships modeling, 257, 267
- ArcHSI, *see* Arc Habitat Suitability Index
- ARM, *see* Adaptive Resource Management
- Artificial neural network (ANN)
 - habitat-relationships modeling, 257, 267
 - multispecies habitat modeling, 237

B

- Bachma's sparrow (*Aimophila aestivalis*), 236
- Bald eagle (*Haliaeetus leucocephalus*), 563
- Balsam fir (*Abies balsamifera*), 485
- Band-tailed pigeon (*Columba fasciata*), 658
- Barn swallow (*Hirundo rustica*), 659
- Barred owl (*Strix varia*), 580
- Bayesian belief network (BBN)
 - habitat-relationships modeling, 258, 269
 - population viability evaluation, 234–235
- BBN, *see* Bayesian belief network

- Beech, 370
- Bell's sage sparrow (*Amphispiza belli*), 240
- BEST, *see* Biodiversity Expert System Tool
- Bewick's wren (*Thyomanes bewickii*), 658
- Bi-criteria species covering problem, reserve selection modeling, 135–139
- BIOCLIM program, 367
- Biodiversity conservation
 - error effects
 - analysis, 103–105
 - implications, 105–107
 - overview, 94–95, 101–103
 - geographic approaches
 - context setting, 87–88
 - data types, 91–92
 - overview, 92–94
 - species criteria, 88–91
 - overview, 85–86
 - research needs
 - data availability and quality, 107
 - ecological process incorporation, 111
 - extended geographic approaches, 108–110
 - inferential basis for conservation planning, 108
 - practitioner accessibility, 111–113
 - scale effects
 - analysis, 96–100
 - extent, 95
 - grain, 95
 - implications, 100–101
 - overview, 94–96
- Biodiversity Expert System Tool (BEST), habitat-relationships modeling, 258, 269
- BIOMAPPER program, 367
- Bird conservation, *see also individual species*
 - Five Elements process
 - assumptions, 616
 - complex species-habitat relationships, 615
 - data constraints and limitations, 615–616
 - flexibility and adaptability, 617
 - model fit, 617
 - projections, 618
 - spatial scalability, 617–618
 - time and cost, 617–619
 - uncertainty, 616–617
 - geographic information system

- Bird conservation (*Continued*)
 - habitat assessment and landscape characterization, 596–598
 - Habitat Suitability Index models, 603–606
 - initiatives in United states, 593–596
 - linking bird distribution and abundance with habitat assessments in Bird Conservation Regions
 - database models, 599
 - geographic information system-based habitat suitability models, 603–606
 - statistical models, 607–614
 - urbanization response modeling
 - avian species richness coupling with relative abundance models, 633–634, 637–645
 - avian surveys, 629–630, 635–637
 - findings, 648–649
 - future land cover prediction, 634–635
 - land-cover change modeling, 630–633
 - limitations, 651–653
 - overview, 625–627
 - policy and management implications, 649–651
 - prospects, 653–654
 - study area and sites, 627–629
 - Urbansim, 655
 - BIRDHAB, habitat-relationships modeling, 258, 267, 278
 - Bison (*Bos bison*), 127
 - Black bear (*Ursus americana*), 65, 231, 380
 - Black-capped chickadee (*Poecile atricapillus*), 658
 - Black-capped vireo (*Vireo atricapillus*), 191
 - Black-footed ferret (*Mustela nigripes*), 167
 - Black-headed grosbeak (*Pheucticus melanocephalus*), 658
 - Black spruce (*Picea mariana*), 484
 - Black tern (*Cblidonia niger*), 534
 - Black-throated gray warbler (*Dendroica nigrescens*), 658
 - Black-throated sparrow (*Amphispiza bilineata*), 517
 - Blue-winged teal (*Anas discors*), 540
 - Bobcat (*Lynx rufus*), 380
 - Bobolink (*Dolichonyx orzivorus*), 211
 - Bobwhite (*Colinus virginianus*), 373, 380
 - BOREAL, habitat-relationships modeling, 258, 267
 - Brewer's blackbird (*Euphagus cyanocephalus*), 517, 659
 - Brewer's sparrow (*Spizella breweri*), 516
 - Brown creeper (*Certhia americana*), 298, 332, 485, 658
 - Brown trout (*Salmo trutta*), 69
 - Brown-headed cowbird (*Molothrus ater*), 609, 659
 - Buffalo (*Bubalus bubalis*), 458–459
 - Bull trout (*Salvenus confluentus*), 563
 - Burrowing owl (*Athene cunicularia*), 167, 600
 - Bushtit (*Psaltriparus minimus*), 658
- ## C
- California gnatcatcher (*Polioptila c. californica*), 461
 - California gull (*Larus californicus*), 534
 - California Wildlife Habitat Relationships (CWHR), habitat-relationships modeling, 258, 267, 278
 - Canada lynx (*Lynx canadensis*), 289, 563
 - Capture/mark/recapture (CMR), 68
 - CART, *see* Classification and regression tree
 - Cascades Habitat Conservation Plan, *see* Plum Creek Central Cascades Habitat Conservation Plan
 - Cassin's vireo (*Vireo cassinii*), 658
 - Cedar, 370
 - Cedar waxwing (*Bombycilla cedrorum*), 658
 - Cerulean warbler (*Dendroica cerulea*), 215, 371, 380
 - Cheatgrass (*Bromus tectorum*), 170, 403, 514, 526
 - Chestnut-backed chickadee (*Poecile ruescens*), 658
 - Chestnut-collared longspur (*Calcarius ornatus*), 161
 - Chisel-toothed kangaroo rat (*Dipodomys microps*), 516
 - Classification and regression tree (CART), 607–608
 - CMR, *see* Capture/mark/recapture
 - Coarse filter
 - versus fine filter approach, 663–664
 - mixed coarse and fine filter approach, 61–62
 - multispecies conservation planning, 56–58
 - Coho salmon (*Oncorhynchus kisutch*), 236
 - Common sagebrush lizard (*Sceloporus graciosus*), 517
 - Common yellowthroat (*Geothlypis trichas*), 658
 - Complementarity, species representation, 89
 - Complex models, utility, 6–7
 - CompPATS, *see* Computerized Project Analysis and Tracking System

- Computerized Project Analysis and Tracking System (CompPATS), habitat-relationships modeling, 258, 265–266, 269
- Conservation efficiency, 203
- Cowbird (*Molothrus ater*), 159
- CWHR, *see* California Wildlife Habitat Relationships
- D**
- Dark kangaroo mouse (*Microdipodops megacephalus*), 517
- Dark-eyed junco (*Junco hyemalis*), 658
- Decision-support model, 2
- Desert horned lizard (*Pbrynosoma platyrhinos*), 516
- Desert spiny lizard (*Sceloporus magister*), 516
- DISPATCH, 323
- DLMP models, *see* Dynamic landscape metapopulation models
- DOMAIN program, 367
- Douglas fir (*Pseudotsuga menziesii*), 567
- Downy woodpecker (*Picoides pubescens*), 658
- Dynamic landscape metapopulation (DLMP) models
- case studies
 - key findings, 487–490
 - modeling approach and scenarios, 485–487
 - Ontario White River management area, 484–485
 - species, 485
 - development steps, 480–481
 - feasibility, 490–491
 - habitat supply model comparison, 491
 - indicator species, 477–479
 - interpretation, 482–484
 - limitations, 492–494
 - overview, 476–477
 - prospects, 494
 - software, 481
- Dynamic species covering problem, reserve selection modeling, 140
- E**
- EAM, *see* Effective Area Model
- Earth Systems Research Institute (ESRI), 290
- Eastern white cedar (*Thuja occidentalis*), 485
- Ecological engineer species, multispecies conservation planning, 61
- Ecosystem Management Decision Support (EMDS), habitat-relationships modeling, 259, 267
- Ecosystem Management Research Institute (EMRI), 172
- Effective Area Model (EAM), habitat-relationships modeling, 258, 268, 278
- Effective population size, population viability analysis, 37–38
- Elk (*Cervus elaphus*), 127, 167, 239, 416, 437–438, 576
- EMBYR, 324
- EMDS, *see* Ecosystem Management Decision Support
- EMP, *see* Expected minimum population size
- Empirical observation, species occupancy, 91
- EMRI, *see* Ecosystem Management Research Institute
- Endangered Species Act (ESA), 51, 153, 155, 163, 561
- Engelman spruce (*Picea engelmanni*), 568
- Environmental Quality Incentives Program (EQIP), 173
- EQIP, *see* Environmental Quality Incentives Program
- ESA, *see* Endangered Species Act
- ESRI, *see* Earth Systems Research Institute
- European starling (*Sturnus vulgaris*), 659
- Evaluation of models, *see* Validation, landscape models
- Expected minimum population size (EMP), dynamic landscape metapopulation modeling, 482
- Expert judgment, biodiversity conservation, 91
- Expert systems, habitat-relationships modeling, 259, 268
- Explanatory model, 2
- F**
- Federal lands
- legislation, 51
 - multispecies conservation planning, 73–76
- Ferruginous hawk (*Buteo regalis*), 167, 517
- FIA, *see* Forest Inventory and Analysis
- Fine filter, multispecies conservation planning
- versus coarse filter, 663–664
 - ecological engineer species, 61
 - focal species, 59–60
 - keystone species, 60
 - link species, 60
 - mixed coarse and fine filter approach, 61–62
 - strong interactor species, 60
 - threatened species, 58–59
 - umbrella species, 60
- Fir, understory vegetation modeling, 403–405

FIRESCAPE, 323
 Flammulated owl (*Otus flammeolus*), 574
 Focal species
 classification, 59–60
 fine filter approach for multispecies
 conservation planning, 59–60
 identification, 71–73
 strategic habitat conservation, 205–206, 223
 Forest Inventory and Analysis (FIA), 74, 289, 375, 377, 605–606
 Forest landscape models, *see also* LANDIS;
 Landscape Management System
 landscape process models, 323–324
 overview, 321–322
 program comparison, 359–360
 succession pathway models, 324
 types, 424–426
 validation, *see* Validation, landscape models
 vital attribute models, 325
 Forest Vegetation Simulator (FVS), 225, 398–399, 425
 FORHAB, habitat-relationships modeling, 259, 269
 FRAGSTATS program, 225, 598
 Franklin's gull (*Larus pipixcan*), 534
 FREIGHTS model, 239
 Fuzzy logic, habitat quality indexing, 231
 FVS, *see* Forest Vegetation Simulator

G

Gadwall (*Anas strepera*), 540
 Gap Analysis Program (GAP), 309, 596
 Gap models, ecosystem dynamics models, 397–398
 GAP, *see* Gap Analysis Program
 General Land Office (GLO), vegetation maps, 292–294, 296
 Generalized linear model (GLM), 607
 Genetic monitoring, multispecies conservation planning, 68–69
 Geographic information system (GIS)
 bird conservation
 habitat assessment and landscape
 characterization, 596–598
 Habitat Suitability Index models, 603–606
 habitat modeling
 analysis structure, 288–290
 classification systems
 attribute variability, 300–301
 language, 296–300
 data abundance, 290–292
 large landscape habitat studies, 225, 227, 230, 240–241

Geographic information system (GIS) (*Continued*)
 spatial scale considerations, 302–305
 technological considerations, 308–312
 temporal scale considerations, 305–308
 vegetation change analysis
 boundary inaccuracy limitation, 294–295
 scale-related limitations, 292–294
 underrepresentation of vegetation
 limitation, 296
 private land conservation planning, 170
 Glanville fritillary butterfly (*Melitaea cixia*), 457
 GLM, *see* Generalized linear model
 GLO, *see* General Land Office
 GNN, *see* Gradient-nearest-neighbor
 Golden-crowned kinglet (*Regulus satrapa*), 658
 Gradient-nearest-neighbor (GNN), understory
 vegetation modeling, 401–402
 Grand fir (*Abies grandis*), 568
 GRASS, 598
 Gray flycatcher (*Empidonax wrightii*), 517
 Gray squirrel (*Sciurus carolinensis*), 380
 Gray wolf (*Canis lupus*), 127, 563
 Great Basin collar lizard (*Crotaphytus
 bicolor*), 516
 Great Basin Ecoregion, 514–521
 Great Basin spadefoot (*Spea intermontana*), 517
 Green-tailed towhee (*Pipilo chlorurus*), 517
 Grizzly bear (*Ursos arctos*), 126, 153, 563
 Groundsnake (*Sonora semiannulata*), 516

H

HABIT@, habitat-relationships modeling, 259, 269
 Habitat-based species viability (HBSV), habitat-relationships modeling, 259, 267
 Habitat Capacity Index (HCI), habitat-relationships modeling, 260, 267
 Habitat Conservation Plan, *see* Plum Creek Central
 Cascades Habitat Conservation Plan
 Habitat Effectiveness Index (HEI), habitat-relationships modeling, 260, 267
 Habitat Evaluation Procedure (HEP), 229, 368
 Habitat networks
 benefits, 502
 case studies
 Great Basin Ecoregion, 514–521
 Interior Columbia Basin Ecosystem Project, 509–513
 overview, 508–509
 conceptual basis, 502–504
 conservation objectives, 506–507
 definition, 501
 development steps, 504–506

- Habitat networks (*Continued*)
 management
 direct links, 522–523
 integration within and among species groups, 521–522
 prospects
 implementation through adaptive management, 525–526
 knowledge gaps, 524–525
 reconciliation of network strategies with other objectives, 523–524
 spatial and temporal basis, 507–508
 HABITAT program, 367
 Habitat Quality (HQ), habitat-relationships modeling, 260, 268
 Habitat quality indexing (HQI), 229–231
 Habitat-relationships modeling
 applications
 cluster 1 frameworks, 273–274
 cluster 2 frameworks, 274–275
 cluster 3 frameworks, 275
 cluster 4 frameworks, 276
 cluster 5 frameworks, 276
 cluster 6 frameworks, 276
 cluster 7 frameworks, 276
 frameworks
 description, 255–256, 266–272
 identification and rating, 252–266
 overview, 251–252
 prospects, 277–278
 Habitat Suitability Index (HSI)
 equations, 379–380
 geographic information system-based bird conservation, 603–606
 habitat-relationships modeling, 260
 Hoosier National Forest case study, 382–385
 input data layers, 375–379
 Landscape HSImodels software, 380–382
 large landscape habitat quantification, 229, 230, 234, 237–238
 model development
 data sources, 371–372
 geographic extent, 370
 spatial grain and extent, 370–371
 suitability functions
 area sensitivity, 372
 distance, 372–373
 edge effects, 373
 habitat type and structure, 372
 landscape composition, 371–373
 overview of models, 368–369
 validation of models, 432
 HABS, *see* Hierarchical All-Bird Strategy
 HABSCAPES, habitat-relationships modeling, 259, 266, 269
 HABSIM, habitat-relationships modeling, 259, 265, 268
 Hairy woodpecker (*Picoides villosus*), 658
 Hammond's flycatcher (*Empidonax hammondi*), 658
 HARVEST, 323, 426–427
 HBSV, *see* Habitat-based species viability
 HCI, *see* Habitat Capacity Index
 HEI, *see* Habitat Effectiveness Index
 Henslow's sparrow (*Ammodramus henslowii*), 380
 HEP, *see* Habitat Evaluation Procedure
 Hermit thrush (*Catbarus guttatus*), 658
 Hickory, 370
 Hierarchical All-Bird Strategy (HABS), 599–602, 615–617
 HNF, *see* Hoosier National Forest
 Hooded warbler (*Wilsonia citrina*), 333, 380
 Hoosier National Forest (HNF)
 decision framework in model selection, 670–672
 Habitat Suitability Index case study, 382–385, 605
 Hotspot analysis, biodiversity conservation, 93
 House finch (*Carpodacus mexicanus*), 659
 House sparrow (*Passer domesticus*), 659
 HQ, *see* Habitat Quality
 HQI, *see* Habitat quality indexing
 HQI program, habitat-relationships modeling, 260, 268
 HQI+ program, habitat-relationships modeling, 260
 HSI, *see* Habitat Suitability Index
 Human-wildlife interactions, *see* People-wildlife interactions
 Hunting, 125–126
 Hutton's vireo (*Vireo buttoni*), 659
 I
 IAN, 598
 IBMs, *see* Individual-based models
 IBR model, *see* Isolation-by-resistance model
 ICBEMP, *see* Interior Columbia Basin Ecosystem Project
 IFM, *see* Incidence function model
 IFOV, *see* Instantaneous Field of View
 Incentives, private land conservation, 163–165
 Incidence function model (IFM), 457
 Indiana bat (*Myotis sodalis*), 380

- Individual-based models (IBMs), large-scale population viability analysis, 463
- Instantaneous Field of View (IFOV), 308–309
- Interior Columbia Basin Ecosystem Project (ICBEMP), 509–513
- Irreplaceability, land, 90
- Isolation-by-resistance (IBR) model, 65
- J**
- Jack pine (*Pinus banksiana*), 127, 191, 331, 484
- Joint ventures, bird conservation, 584–595
- K**
- K-nearest-neighbor (KNN), understory vegetation modeling, 401
- Kentucky warbler (*Oporornis formosus*), 609–613
- Keystone species
- dynamic landscape metapopulation models, 477
 - multispecies conservation planning, 60
- Killdeer (*Charadrius vociferus*), 658
- King rail (*Rallus elegans*), 207
- Kirtland's warbler (*Dendroica kirtlandii*), 127, 191, 193
- Kit fox (*Vulpes macrotis*), 517
- KNN, *see* K-nearest-neighbor
- L**
- LAFS, 324
- Lake sturgeon (*Acipenser fulvescens*), 207
- Land Use Evolution and Impact Assessment Model (LEAM), habitat-relationships modeling, 261, 267
- Land-cover change (LCC) modeling, 630–633, 647
- LANDFIRE, *see* Landscape Fire and Resource Management Planning Tools Project
- LANDIS
- dynamic landscape metapopulation modeling, 481
 - Landscape Management System
 - comparison, 359
 - natural disturbance, 327–328
 - operational design, 328–329
 - overall design and structure, 325–326
 - succession, 326–327
 - timber harvest and fuel treatment, 328
 - versions, 334
 - wildlife model integration
 - advantages and limitations, 333–335
 - approaches, 329–333
 - Landscape connectivity, addressing for multispecies conservation planning, 54, 64–65
 - A Landscape Cumulative Effects Simulator (ALCES), habitat-relationships modeling, 257, 267
 - Landscape Evaluation Effects of Management Effects on Timber and Habitat (LEEMATH), habitat-relationships modeling, 239, 261, 265, 267
 - Landscape Fire and Resource Management Planning Tools Project (LANDFIRE), 597
 - Landscape HSI, habitat-relationships modeling, 261, 268
 - Landscape Management System (LMS)
 - comparison with other programs
 - LANDIS, 359
 - Spatial Woodstock, 360
 - decision analysis
 - alternate landscape plan development, 355–358
 - grouping stands, 355
 - overview, 354
 - stand-specific prescriptions from groups, 357, 359
 - geographic area of study and application, 344–345
 - habitat-relationships modeling, 261, 267
 - management application, 351–354
 - modules, 341–343
 - output
 - criteria for sustainable forestry, 349–350
 - inventory information, 348
 - summarized information, 348
 - visualization, 348–349
 - portfolio creation, 345–346
 - prospects, 360–362
 - scientific basis
 - complexity and systems theory, 341–342
 - decision analysis, 342
 - hierarchies and top-down planning, 342
 - stand and landscape dynamics, 340–341
 - trade-offs, 341
 - stand treating and projecting, 346–348
 - Toggle program, 356–357
 - Landscape models, *see also* Forest landscape models
 - Adaptive Resource Management, 11–12
 - decision framework in model selection
 - case studies, 671–672
 - coarse versus fine filter, 663–664
 - conceptual versus statistical models, 667

- Landscape models (*Continued*)
- existing versus new data collection, 666
 - habitat, abundance, or viability, 665–666
 - optimization versus choosing from
 - alternatives, 669–671
 - overview, 661
 - reserve selection versus land management
 - planning, 662–663
 - single species versus multiple species, 663
 - species representation versus diversity,
 - 664–665
 - stationary versus dynamic approaches,
 - 667–669
 - overview, 1–2
 - philosophy of modeling, 3–10
 - uncertainty, 12–16
 - unreliable model avoidance, 10–11
 - uses, 2
 - validation, *see* Validation, landscape models
 - viability a general management metric, 16–20
 - visualization, 20–26
- LANDSIM, 325
- LANDSUM, 324, 426–427
- Large landscape habitat quantification
- comparison of landscapes
 - spatial comparisons, 238–239
 - temporal comparisons, 239–240
 - complexity gradient of models, 226–229
 - geographic information system
 - analysis structure, 288–290
 - classification systems
 - attribute variability, 300–301
 - language, 296–300
 - data abundance, 290–292
 - examples, 225, 227, 230, 240–241
 - spatial scale considerations, 302–305
 - technological considerations, 308–312
 - temporal scale considerations, 305–308
 - vegetation change analysis
 - boundary inaccuracy limitation, 294–295
 - scale-related limitations, 292–294
 - underrepresentation of vegetation
 - limitation, 296
- Habitat Suitability Index modeling, *see* Habitat Suitability Index
- model evaluation, 240–241
 - multispecies approaches, 236–238
 - occurrence probability modeling, 231–233
 - overview, 225–226
 - population density estimation, 233–234
 - population viability evaluation, 234–236
 - prospects, 242
- Large landscape habitat quantification
- (*Continued*)
 - quality indexing, 229–231
- Lark bunting (*Calamospiza melanocorys*), 161
- Lark sparrow (*Chodestes grammacus*), 517
- Lattice models, large-scale population viability analysis, 458–460
- LCC modeling, *see* Land-cover change modeling
- LEAM, *see* Land Use Evolution and Impact Assessment Model
- LEEMATH, *see* Landscape Evaluation Effects of Management Effects on Timber and Habitat
- Lesser prairie-chicken (*Tympanuchus pallidicinctus*), 601–602
- LiDAR, *see* Light Detection and Ranging
- Light Detection and Ranging (LiDAR), Habitat Suitability Index input, 378–379
- Link species, multispecies conservation planning, 60
- Little pocket mouse (*Perognathus longimembris*), 517
- LMS, *see* Landscape Management System
- Lodgepole pine (*Pinus contorta*), 568
- Loggerhead shrike (*Lanus ludovicianus*), 167, 517
- Long-billed curlew (*Numenius americanus*), 161, 167
- Long-nosed leopard lizard (*Gambelia wislizeni*), 516
- Long-nosed snake (*Rhinocbeilus lecontei*), 516
- ## M
- MacGillivray's warbler (*Oporornis tolmei*), 659
- Magpie goose (*Anseranas semipalmata*), 460
- Mallard (*Anas platyrhynchos*), 540
- Management-oriented model, 9–10
- Maple, 370
- Marbled godwit (*Limosa fedoa*), 210, 533, 545–546, 549–551
- Marbled murrelet (*Brachyramphus mamoratus*), 563
- MARS, *see* Multivariate adaptive regression splines
- Marten (*Martes americana*), 239
- MARXAN program, 97
- Matrix models, population viability analysis, 35
- MAXENT program, 64
- Maximum expected species covering problem, reserve selection modeling, 139
- Maximum species covering problem, reserve selection modeling, 131–135
- MBCE, *see* Migratory Bird Conservation Fund
- McCown's longspur (*Calcarius mccownii*), 161

- MDNR, *see* Michigan Department of Natural Resources
- Measurement scale, geographic information system, 303–305
- Merriam's kangaroo rat (*Dipodomys merriami*), 516
- Merriam's shrew (*Sorex merriami*), 517
- Mesofilter, multispecies conservation planning, 58
- Metadata, geographic information system, 290
- Metapopulation models
 - dynamic landscape metapopulation models, *see* Dynamic landscape metapopulation models
 - large-scale population viability analysis, 460–462, 475
 - population viability analysis, 36
- Metapopulation size problem, reserve design modeling, 143
- Michigan Department of Natural Resources (MDNR)
 - map boundary limitations, 294–295
 - vegetation classification, 297–298
- Migratory Bird Conservation Fund (MBCF), 535
- Missouri Forest Ecosystem Project (MOFEP), 441
- MOFEP, *see* Missouri Forest Ecosystem Project
- Moose (*Alces alces*), 207
- Most-similar-neighbor (MSN), understory vegetation modeling, 401
- Mountain hemlock (*Tsuga mertensiana*), 567
- Mountain plover (*Charadrius montanus*), 161, 167
- Movement, landscape connectivity in multispecies conservation planning, 54
- MSN, *see* Most-similar-neighbor
- Mule deer (*Odocoileus bemonius*), 167, 416
- Multispecies conservation planning
 - allometric scaling relationships, 53
 - demography, 54
 - federal public lands, 73–76
 - filtering approaches
 - coarse filter, 56–58
 - fine filter
 - threatened species, 58–59
 - focal species, 59–60
 - umbrella species, 60
 - keystone species, 60
 - strong interactor species, 60
 - link species, 60
 - ecological engineer species, 61
 - mesofilter, 58
 - mixed coarse and fine filter approach, 61–62
 - overview, 55–56
- Multispecies conservation planning (*Continued*)
 - focal species identification, 71–73
 - genetic monitoring, 68–69
 - landscape connectivity addressing for multiple species, 64–65
 - movement and landscape connectivity, 54
 - The Nature Conservancy strategy, 69–70
 - niche theory, 52–53
 - persistence likelihood, maximization for multiple species, 67–68
 - probability surfaces for species occurrences, 62–64
 - reserve selection algorithms, 65–67
 - versus single species, 663
 - trophic structure, 53
- Multivariate adaptive regression splines (MARS), 608
- ## N
- NABCI, *see* North American Bird Conservation Initiative
- National Elevation Dataset (NED), 598
- National Forest Management Act (NFMA), 51, 59, 73
- National Hydrography Dataset (NHD), 598
- National Land Cover Dataset (NLCD), 596–597
- National Vegetation Classification, 299
- National Wetlands Inventory (NWI), 294–295, 539–540
- The Nature Conservancy (TNC), multispecies conservation planning strategy, 69–70
- NAWCA, *see* North American Wetlands Conservation Act
- NAWMP, *see* North American Waterfowl Management Plan
- NED, *see* National Elevation Dataset
- NFMA, *see* National Forest Management Act
- NHD, *see* National Hydrography Dataset
- Niche theory, multispecies conservation planning, 52–53
- Nightsnake (*Hypsiglena torquata*), 517
- NLCD, *see* National Land Cover Dataset
- Noble fir (*Abies procera*), 567
- North American Bird Conservation Initiative (NABCI), 594
- North American Waterfowl Management Plan (NAWMP), 594
- North American Wetlands Conservation Act (NAWCA), 535
- Norther rough-winged swallow (*Stelgidopteryx serripennis*), 659
- Northern flicker (*Colaptes auratus*), 659

- Northern goshawk (*Acciper gentilis*), 191
 Northern grasshopper mouse (*Onychomys leucogaster*), 517
 Northern harrier (*Circus cyaneus*), 517
 Northern long-eared bat (*Myotis septentrionalis*), 380
 Northern pintail (*Anas acuta*), 540
 Northern shoveler (*Anas clypeata*), 540
 Northern spotted owl (*Strix occidentalis*), 59, 124, 191, 235, 506, 561-588
 Nuthatch (*Sitta europaea*), 233
 NWI, *see* National Wetlands Inventory
- O**
- Oak, 370, 429-430
 Object-oriented programming (OOP), habitat quality indexing, 231
 Observational scale, geographic information system, 302-303
 Occam's razor, 6
 Occupancy models, large-scale population viability analysis, 455, 457-458
 Olive-sided flycatcher (*Contopus cooperi*), 659
 ONFIRE, 323
 OOP, *see* Object-oriented programming
 Optimization analysis, biodiversity conservation, 94
 OPTIONS program, futuring of wildlife habitat, 574-576
 Orange-crowed warbler (*Vermivora celata*), 659
 Ord's kangaroo rat (*Dipodomys ordii*), 517
 Ornate box turtle (*Terrapene ornata*), 212
 Ovenbird (*Seiurus aurocapillus*), 240, 332, 380
- P**
- Pacific fisher (*Martes pennanti*), 574
 Pacific silver fir (*Abies amabilis*), 567
 Pacific-slope flycatcher (*Empidonax difficilis*), 634, 640-641, 646, 659
 Parameters, model
 estimation, 10, 14-15
 dependence on too many, 10
 Partners in Flight (PIF), 594-595, 614-615
 Patch models, population viability analysis, 36
 PATCH, *see* Program to Assist in Tracking Critical Habitat
 PATREC, *see* Pattern Recognition
 Pattern Recognition (PATREC)
 habitat-relationships modeling, 262, 267
 population density estimation, 234
 People-wildlife interactions
 People-wildlife interactions (*Continued*)
 amenity migration, 127-129
 recreation, 124-126
 restoration, 126-127, 148
 Peregrine falcon (*Falco peregrinus*), 563
 Persistence likelihood, maximization for multiple species, 67-68
 Pied-billed grebe (*Podilymbus podiceps*), 534
 PIF, *see* Partners in Flight
 Pig, (*sus scrofa*), 458
 Pileated woodpecker (*Dryocopus pileatus*), 574
 Pine siskin (*Carduelis pinus*), 659
 Pine warbler (*Dendroica pinus*), 333, 380
 Pinyon-juniper woodlands, 514
 Playa Lakes Joint Venture (PLJV), 599-600, 602
 PLJV, *see* Playa Lakes Joint Venture
 Plum Creek Central Cascades Habitat Conservation Plan
 adaptive management and monitoring, 582-584
 forest stand classification, 570-571
 futuring of wildlife habitat, 574-577
 Habitat Conservation Plan process, 563-566
 land ownership and area, 566-568
 lifeforms and habitat analysis, 571-574
 multispecies approach, 568-570
 northern spotted owl
 centers, 562
 resource selection function analysis, 578-579
 resource selection probability function, 580-581
 policy and management implications, 584-587
 prospects, 587-588
 threatened species, 563
 Point Specific Estimator (PSE), habitat-relationships modeling, 262, 268
 Ponderosa pine (*Pinus ponderosa*), 568
 Population viability analysis (PVA),
 see also Viability
 decision theory, 40
 demographic models
 matrix models, 35
 metapopulation models, 36
 spatially explicit population models, 36
 stochastic population viability analysis models, 35-36
 effective population size, 37-38
 large landscape applications
 connectivity, 453
 habitat quantification, 234-236
 model selection and parameterization, 464
 model types and applications
 individual-based models, 463

- Population viability analysis (PVA) (*Continued*)
- lattice models, 458–460
 - metapopulation models, 460–462
 - occupancy models, 455, 457–458
 - overview, 456
 - prospects, 465–466
 - rationale, 449–450
 - spatial extent, 451–452
 - spatial heterogeneity, 453–454
 - spatial resolution, 452–453
 - temporal dynamics, 454–455
 - model selection factors, 38–39
 - optimization framework applications, 41–42
 - relationship to other methods, 450–451
 - scope, 33
 - stochastic patch occupancy models, 37
 - trend analysis, 37
 - uncertainty, 42–43
- PPR, *see* Prairie Pothole Region
- Prairie dog (*Cynomys*), 143, 159, 167, 600
- Prairie falcon (*Falco mexicanus*), 517
- Prairie Pothole Region (PPR)
- conservation efforts, 534–535
 - conservation planning approach
 - data availability and quality, 539–540
 - ducks, 540
 - model development and integration, 536–539
 - realities, 535–536
 - waterbirds and shorebirds, 540–542
 - overview, 533–534
 - policy and management implications of
 - conservation, 554–555
 - prospects for conservation, 555–556
 - species abundance and distribution models
 - conservation applications, 551–554
 - marbled godwit
 - conceptual model, 546
 - outcomes, 549–551
 - statistical models, 545–546
 - sora, 543–545, 548–549
 - waterfowl, 542–543, 547–548
- Prairie warbler (*Dendroica discolor*), 333, 380
- Predicted occurrence, species in biodiversity
 - conservation, 92
- Predictive model, 2, 5
- Primary viability, 18
- Private lands
- abundance, 153
 - conservation planning challenges
 - funding, 155–156
 - landowner engagement, 156–157
 - mixed ownership, 159–160
- Private lands (*Continued*)
- overview, 154
 - property rights, 154–155
 - proprietary information, 157–158
 - solutions, 160–165
 - species management focus, 158–159
 - ecosystem diversity approach for conservation
 - planning, 160–163
 - incentives for conservation planning, 163–165
 - modeling
 - forestlands, 166–167
 - Thunder Basin Grassland Prairie Ecosystem Association, 167–173
- PROC CLUSTER, habitat-relationships
 - modeling, 256
- PROC DISTANCE, habitat-relationships
 - modeling, 256
- Program to Assist in Tracking Critical Habitat (PATCH), habitat-relationships
 - modeling, 261, 267
- Pronghorn (*Antilocapra americana*), 517
- Pronghorn antelope (*Antilocapra americana*), 167
- PSE, *see* Point Specific Estimator
- Public lands, *see* Federal lands
- Purple finch (*Carpodacus purpureus*), 659
- Purpose, defining for model, 4–5
- PVA, *see* Population viability analysis
- Pygmy rabbit (*Brachylagus idahoensis*), 516
- ## R
- RAMAS LANDIS, 331
- RAMAS Landscape
- applications, 239–240
 - habitat-relationships modeling, 262, 268
 - limitations, 492–494
- RAMAS GIS module
- dynamic landscape metapopulation modeling, 481, 485, 490
 - LANDIS integration, 332
- Real estate investment trust (REIT), 166
- Recreation, 124–126
- Red-backed salamander (*Plethodon cinereus*), 485
- Red-backed vole (*Clethrionomys gapperi*), 485
- Red bat (*Lasturus borealis*), 380
- Red-breasted nuthatch (*Sitta canadensis*), 659
- Red-cockaded woodpecker (*Picoides borealis*), 157
- Red crossbill (*Loxia curvirostra*), 659
- Red pine (*Pinus resinosa*), 331
- Red-winged blackbird (*Agelaius phoeniceus*), 659
- Redundancy, species incidence, 90

- Region specificity, models, 8–9
- REIT, *see* Real estate investment trust
- Reserve-based modeling
- advantages and limitations, 146–147
 - overview, 129–131
 - prospects, 147–148
 - reserve design models
 - population dynamics incorporation
 - metapopulation size problem, 143
 - overview, 142–143
 - safe-minimum-standard problem, 144–145
 - surviving populations problem, 145–146
 - reserve compactness problem, 142
 - reserve connectivity problem, 141–142
 - reserve proximity problem, 141
 - reserve selection models
 - bi-criteria species covering problem, 135–139
 - decision framework in model selection, 662–663, 672
 - dynamic species covering problem, 140
 - maximum expected species covering problem, 139
 - maximum species covering problem, 131–135
 - multispecies conservation planning, 65–67
- Resource selection function (RSF), 578–579
- Resource selection probability function (RSPF), 580–581
- Restoration, habitats, 126–127, 148
- Robustness, biodiversity conservation, 90
- Rock pigeon (*Columba livia*), 659
- Rock squirrel (*Spermophilus variegatus*), 518
- Rock wren (*Salpinctes obsoletus*), 518
- RPF, *see* Resource selection probability function
- RSF, *see* Resource selection function
- Ruffed grouse (*Bonasa umbellus*), 380
- Rufous hummingbird (*Selasphorus rufus*), 659
- S**
- Safe-minimum-standard problem, reserve design modeling, 144–145
- Sage grouse (*Centrocercus urophasianus*), 156, 236, 403, 516
- Sage sparrow (*Amphispiza belli*), 516
- Sage thrasher (*Oreoscoptes montanus*), 516
- Sagebrush (*Artemisia*), 157, 518
- Sagebrush vole (*Lemmingscurtus*), 517
- Savannah sparrow (*Passerculus sandwichensis*), 659
- Secondary viability, 18
- SEM, *see* Spatially explicit model
- SEPMs, *see* Spatially explicit population models
- SESI, *see* Spatially Explicit Species Index
- Sharp-tailed grouse (*Tympanuchus phastanellus*), 240, 310, 331, 462
- SHC, *see* Strategic habitat conservation
- SHM, *see* Species-habitat matrices
- Short-eared owl (*Asio flammeus*), 161, 517
- SIMFOR, habitat-relationships modeling, 262, 268
- SIMPPLLE, 324, 426–427
- SNBN, *see* Spatially Neutral Bayesian Model
- Snowshoe hare (*Lepus americanus*), 301
- Song sparrow (*Melospiza melodia*), 659
- Sora (*Porzana carolina*), 210, 533–534, 543–545, 548–549
- Southern redback salamander (*Plethodon serratus*), 280
- Spatial optimization, habitat-relationships modeling, 263, 268
- Spatial Woodstock, Landscape Management System comparison, 360
- Spatially Explicit Species Index (SESI), habitat-relationships modeling, 262, 268
- Spatially explicit model (SEM), strategic habitat conservation, 212
- Spatially explicit population models (SEPMs), 36
- Spatially Neutral Bayesian Model (SNBN), habitat-relationships modeling, 263, 268
- Species-area relationship (SPPAREA), habitat-relationships modeling, 263, 268
- Species count, biodiversity conservation, 88.89
- Species-habitat matrices (SHM), habitat-relationships modeling, 263, 267, 278
- Species representation, biodiversity conservation, 89
- Species richness, *see* Species count
- Species sorting algorithm (SSA), habitat-relationships modeling, 263, 269
- Spectral-distance error, geographic information system, 309–310
- SPOMs, *see* Stochastic patch occupancy models
- Spotted owl, *see* Northern Spotted owl
- Spotted towhee (*Pipilo maculatus*), 659
- SPPAREA, *see* Species-area relationship
- SSA, *see* Species sorting algorithm
- Steller's jay (*Cyanocitta stelleri*), 659
- Stepwise approximation approach, conservation assessment and planning
 - broad-scale multispecies ranges, 181
 - examples
 - multiscale bird distribution assessment, 182–184
 - openland bird assessment, 189–191

- Stepwise approximation approach, conservation assessment and planning (*Continued*)
- habitat-quality evaluation
 - abundance estimates and temporal change, 187–188
 - refinement with productivity and survivorship estimates, 188–189
 - implementation guidelines, 194–195
 - maps and summary models, 191
 - overview, 179–181
 - rationale, 177–179
 - resolution levels
 - level 1, 181–184
 - level 2, 185–186
 - level 3, 186–187
 - level 4, 187–188
 - level 5, 188–189
 - level 6, 191–192
 - level 7, 192–194
 - species distribution refinement via habitat-gradient relationships, 185–186
 - testing and integrating monitoring into science, 192–194
 - test species habitat or gradient relationships across landscapes, 186–187
- Stochastic patch occupancy models (SPOMs), population viability analysis, 37
- Stochastic population viability analysis models, 35–36
- Strategic habitat conservation (SHC)
- advantages, 220–221
 - biological planning
 - focal species selection, 205–206, 223
 - limiting factor and conservation treatment identification, 208
 - model development and application, 208–211
 - population objective setting, 206–207
 - conservation design
 - biodiversity and species richness maps, 214–216
 - ecosystem assessment, 212
 - habitat objectives, 213
 - priority area designation, 213–214
 - spatially explicit model development, 212
 - evaluation
 - assumption-driven research, 216–218
 - conservation action effects, 218–219
 - outcome-based monitoring, 218
 - population objective progress, 220
 - program and agency accomplishments, 219–220
- Strategic habitat conservation (SHC) (*Continued*)
- forest type matrix, 223
 - overview, 202–204
 - rationale, 201–202
- Striped whipsnake (*Masticophis taeniatus*), 517
- Strong interactor species, multispecies conservation planning, 60
- Subalpine fir (*Abies lasiocarpa*), 568
- Surrogates, biodiversity conservation, 92
- Surviving populations problem, reserve design modeling, 145–146
- Swainson's hawk (*Buteo swainsoni*), 517
- Swainson's thrush (*Catharus ustulatus*), 659
- ## T
- Tamarack (*Larix laricina*), 485
- TBGPEA, *see* Thunder Basin Grassland Prairie Ecosystem Association
- Tertiary viability, 18–19
- Thunder Basin Grassland Prairie Ecosystem Association (TBGPEA), private land conservation modeling, 167–173
- Tiger salamander (*Ambystoma tigrinum*), 522
- Timber rattlesnake (*Crotalus horridus*), 380
- Timber wolf (*Canis lupus lycaon*), 126
- Timberland investment management organization (TIMO), 166
- TIMO, *see* Timberland investment management organization
- TNC, *see* The Nature Conservancy
- Toggle program, *see* Landscape Management System
- Transparency, models, 7–8
- Tree swallow (*Tachycineta bicolor*), 659
- Trembling aspen (*Populus tremuloides*), 484–485
- Trend analysis, population viability analysis, 37
- Turkey (*Melagris gallopavo*), 230, 380
- ## U
- Umbrella species, multispecies conservation planning, 60
- Uncertainty, models
 - hypotheses and model structure, 13–14
 - model user bias, 16
 - parameter estimates, 14–15
 - population viability analysis, 42–43
 - verification, 15–16
- Understory vegetation modeling
 - ecosystem dynamics models, 397–399
 - empirical models
 - abundance data, 395–396

Understory vegetation modeling (*Continued*)
 binary data, 394–395
 data requirements and limitations, 397
 definition, 391–392
 multivariate data, 397
 techniques, 393–394
 fire response, 403–405
 ideal model, 400
 importance, 391
 large landscape predictions, 400–402
 paradigms, 392–393
 process-based models, 392
 prospects, 405–407
 qualitative models, 392, 399–400
 Unreliable models, avoiding, 10–11
 Upland sandpiper (*Bartramia longicauda*),
 161, 167
 Urbanization response modeling, *see* Bird
 conservation

V

Validation, landscape models
 animal location data, 436–440
 autocorrelation, 435–436
 comparison of models, 418
 data, 431–432, 434
 definitions
 calibration, 419
 evaluation, 419
 validation, 419
 verification, 419
 evaluation elements
 application environment, 420, 422
 conceptual design, 422–423
 overview, 421
 quantitative performance, 423–424
 forest vegetation models
 forest structure change validation, 427
 modeled disturbance evaluation,
 42–428
 species change evaluation, 428–430
 types, 424–426
 metrics, 418–419, 430
 model uncertainty, 433–434
 objectives, 418
 overview, 415–418
 prospects, 440–442
 spatial scale and validation, 434–435
 VDDT/TELSA, 426–427
 Verification, models, 15–16
 Vesper sparrow (*Pooecetes gramineus*),
 516

Viability, *see also* Population viability analysis
 definition, 17
 modeling in large landscapes,
 19–20
 primary viability, 18
 secondary viability, 18
 tertiary viability, 18–19
 Violet-green swallow (*Tachycineta thalassina*),
 659
 Visualization, models
 context, 23–24
 human perception issues, 24
 misrepresentation, 24–26
 motion utilization, 22–23
 overview, 20–22
 Vulnerability, biodiversity conservation, 90

W

Warbling vireo (*Vireo gilvus*), 659
 Weighted linear combination (WLC), habitat
 quality indexing, 230–231
 Western burrowing owl (*Athene cubicaria
 hypugaea*), 517
 Western hemlock (*Tsuga heterophylla*),
 567
 Western larch (*Larix occidentalis*), 568
 Western spruce budworm (*Choristoneura
 occidentalis*), 577
 Western tanager (*Piranga ludoviciana*),
 659
 Western white pine (*Pinus monticola*),
 568
 Western wood pewee (*Contopus sordidulus*),
 659
 White birch (*Betula papyrifera*), 485
 White-crowned sparrow (*Zonotrichia
 leucophrys*), 659
 White-headed woodpecker (*Picoides
 albolarvatus*), 522
 White spruce (*Picea glauca*), 485
 White-tailed deer (*Odocoileus virginianus*), 167,
 234, 416, 612
 White-tailed jackrabbit (*Lepus townsendii*),
 517
 WHQ, *see* Wildlife habitat quality
 Wildlife habitat quality (WHQ), habitat-
 relationships modeling, 264, 268
 Willow flycatcher (*Empidonax traillii*),
 659
 Wilson's warbler (*Wilsonia pusilla*), 659
 Winter wren (*Troglodytes troglodytes*), 659
 WLC, *see* Weighted linear combination

Wolverine (*Gulo gulo*), 153
Wood thrush (*Hyocichla mustelina*), 370
Woodland caribou (*Rangifer tarandus*),
239
Worm-eating warbler (*Helmintheros vermivorous*),
609-613
Wyoming ground squirrel (*Spermophilus elegans*
nevadensis), 516

Y

Yellow-breasted chat (*Icterus virens*), 370
Yellow-rumped warbler (*Dendroica coronata*),
634, 640-641, 646, 659

Z

ZONATION program, 63, 65