

## Chapter 22

### Effects of elevated CO<sub>2</sub> and O<sub>3</sub> on aspen clones of varying O<sub>3</sub> sensitivity

B.A. Wustman

*Amicus Therapeutics Inc., Commercialization Center for Innovative Technologies,  
675 US Highway One, North Brunswick, NJ 08902, USA*

E. Oksanen

*University of Kuopio, Department of Ecology and Environmental Science,  
POB 1627, 70211 Kuopio, Finland*

D.F. Karnosky

*Michigan Technological University, School of Forest Resources and Environmental Science,  
1400 Townsend Drive, Houghton, MI 49931-1295, USA*

A. Noormets

*The University of Toledo, Department of EEES, LEES Lab, Mail Stop 604,  
Toledo, OH 43606, USA*

J.G. Isebrands

*Environmental Forestry Consultants, LLC, P.O. Box 54, E7323 Hwy 54,  
New London, WI 54501, USA*

K.S. Pregitzer

*Michigan Technological University, School of Forest Resources and Environmental Science,  
1400 Townsend Drive, Houghton, MI 49931-1295, USA*

G.R. Hendrey

*Brookhaven National Lab, Division of Environmental Biology and Instrumentation,  
1 South Technology Street, Upton, NY 11973, USA*

J. Sober

*Michigan Technological University, School of Forest Resources and Environmental Science,  
1400 Townsend Drive, Houghton, MI 49931-1295, USA*

G.K. Podila\*

*University of Alabama, Department of Biological Sciences, 301 Sparkman Drive,  
WH142, Huntsville, AL 35899, USA*

---

\*Corresponding author.

### Abstract

To determine whether elevated CO<sub>2</sub> reduces or exacerbates the detrimental effects of O<sub>3</sub> on aspen (*Populus tremuloides* Michx.). Aspen clones 216 and 271 (O<sub>3</sub> tolerant), and 259 (O<sub>3</sub> sensitive) were exposed to ambient levels of CO<sub>2</sub> and O<sub>3</sub> or elevated levels of CO<sub>2</sub>, O<sub>3</sub>, or CO<sub>2</sub> + O<sub>3</sub> in the FACTS II (Aspen FACE) experiment, and physiological and molecular responses were measured and compared. Clone 259, the most O<sub>3</sub>-sensitive clone, showed the greatest amount of visible foliar symptoms as well as significant decreases in chlorophyll, carotenoid, starch, and ribulose-1,5-bisphosphate carboxylase/oxygenase (Ru-bisco) concentrations and transcription levels for the Rubisco small subunit. Generally, the constitutive (basic) transcript levels for phenylalanine ammonia-lyase (Pal) and chalcone synthase (Chs) and the average antioxidant activities were lower for the ozone sensitive clone 259 as compared to the more tolerant 216 and 271 clones. A significant decrease in chlorophyll a, b and total (a + b) concentrations in CO<sub>2</sub>, O<sub>3</sub>, and CO<sub>2</sub> + O<sub>3</sub> plants was observed for all clones. Carotenoid concentrations were also significantly lower in all clones; however, Chs transcript levels were not significantly affected, suggesting a possible degradation of carotenoid pigments in O<sub>3</sub>-stressed plants. Antioxidant activities and Pal and 1-aminocyclopropane-1-carboxylic acid (ACC)-oxidase transcript levels showed a general increase in all O<sub>3</sub> treated clones, while remaining low in CO<sub>2</sub> and CO<sub>2</sub> + O<sub>3</sub> plants (although not all differences were significant). Our results suggest that the ascorbate-glutathione and phenylpropanoid pathways were activated under ozone stress and suppressed during exposure to elevated CO<sub>2</sub>. Although CO<sub>2</sub> + O<sub>3</sub> treatment resulted in a slight reduction of O<sub>3</sub>-induced leaf injury, it did not appear to ameliorate all of the harmful affects of O<sub>3</sub> and, in fact, may have contributed to an increase in chloroplast damage in all three aspen clones.

### 1. Introduction

Atmospheric CO<sub>2</sub> is increasing rapidly and is expected to double by the end of the next century (Barnola et al., 1995). Tropospheric ozone (O<sub>3</sub>) is also increasing globally at a rate of 1–2% per year (Chameides et al., 1995) and it is likely that these two gases will have significant impacts in the future on forest tree species and ecosystems (Matyssek and Innes, 1999; Reilly et al., 1999). However, it is not clear at this point whether or not elevated CO<sub>2</sub> reduces or exacerbates the detrimental effects of O<sub>3</sub>. Moreover, leaf age can greatly affect the direction of the CO<sub>2</sub> + O<sub>3</sub> response. McKee et al. (1995) showed no protective effect of elevated CO<sub>2</sub> against O<sub>3</sub> on emerging wheat flag leaves, but strong effects in mature leaves. Experimental results provide evidence for both hypotheses, but neither has been tested under open field conditions, and both could be strongly influenced by the lower boundary layer conductances that characterize enclosures (Polle and Pell, 1999). There is clearly a need to test this in an open-air facility, such as FACE, where artifacts are minimized. The view that elevated CO<sub>2</sub> should reduce the detrimental effects of O<sub>3</sub> is based

primarily on the expected decrease in stomatal conductance in elevated CO<sub>2</sub>, which reduces O<sub>3</sub> flux received by the plant. The increase in substrate availability for repair and detoxification at elevated CO<sub>2</sub> may also reduce the effect of elevated O<sub>3</sub> (Rao et al., 1995).

The alternate hypothesis, that elevated CO<sub>2</sub> will exacerbate the detrimental effects of O<sub>3</sub>, is based primarily on the prediction that reduced photorespiration, decreased Rubisco content, increased availability of CO<sub>2</sub> for photosynthesis, and a decrease in non-photochemical energy dissipation will reduce the need for cellular detoxification of reactive oxygen species (Polle et al., 1993). Thus, the lower antioxidant levels result in reduced tolerance to O<sub>3</sub>. In aspen, there is some evidence of antioxidants playing a key role in O<sub>3</sub> tolerance (Sheng et al., 1997; Noormets et al., 2000) and an apparent increase in O<sub>3</sub> sensitivity with elevated CO<sub>2</sub> (Kull et al., 1996). In aspen softwood cuttings grown in open-top chambers, CO<sub>2</sub> did not compensate for the deleterious effects of elevated O<sub>3</sub> and, in some cases, photosynthetic capacity decreased more than with O<sub>3</sub> alone (Kull et al., 1996). This was particularly true for the O<sub>3</sub>-sensitive clone. Moreover, the tolerant clones sometimes became more sensitive to O<sub>3</sub> with CO<sub>2</sub> enrichment (Kull et al., 1996; Karnosky et al., 1998). Data from various tree species is conflicting on the interacting effects of O<sub>3</sub> and CO<sub>2</sub>. These findings fall into two categories, (1) the elevated CO<sub>2</sub> has some positive effects in ameliorating the negative effects of O<sub>3</sub> (Mortensen, 1995; Kellomaki and Wang, 1997; Dickson et al., 1998; Sehmer et al., 1998; Grams et al., 1999; Utrainen et al., 2000; Broadmeadow et al., 1999) or (2) it increases the negative effects of O<sub>3</sub> (Kull et al., 1996; Karnosky et al., 1998; Niewiadomska et al., 1999).

To better understand the interactive affects of increased CO<sub>2</sub> and O<sub>3</sub> levels, we have measured various leaf antioxidant activities and gene expression patterns of several stress-response pathways for aspen trees growing in a Free Air CO<sub>2</sub> and/or O<sub>3</sub> Enrichment (FACE) experiment. Chlorophyll, carotenoid, and Rubisco contents have also been measured to help determine how the photosynthetic capacities are affected for clones of differing O<sub>3</sub> sensitivity. Results from this open air experiment indicate that elevated CO<sub>2</sub> levels increase O<sub>3</sub> sensitivity in all three clones by suppressing ascorbate-glutathione and phenylpropanoid pathways. An increase in AOS, consequently, may be responsible for the observed decreases in chlorophyll and carotenoid pigments.

## **2. Materials and methods**

### ***2.1. Plant material and fumigation protocols***

The research facility is located in the USDA Forest Service Harshaw farm site in Oneida County, Wisconsin (Karnosky et al., 1999; Dickson et al., 2000).

Three replicate FACE rings were established for each treatment (control (= ambient), elevated O<sub>3</sub>, elevated CO<sub>2</sub>, and elevated O<sub>3</sub> + CO<sub>2</sub>). In elevated-O<sub>3</sub> rings, O<sub>3</sub> concentrations were maintained approximately 1.5 times higher than the ambient concentrations. The mean O<sub>3</sub> concentrations during the 1998 growing season were 36 ppb for ambient air and 56 ppb for elevated O<sub>3</sub> exposures. The monthly AOT40 values ranged from 5.2–9.2 ppm-h (mean 7.0 ppm-h) for elevated O<sub>3</sub> treatments and ranged from 25.9–30.6 ppm-h for the seasonal dose (Table 1). The mean concentrations and total cumulative O<sub>3</sub> exposures over a threshold of 0 ppb (AOT00) and of 40 ppb (AOT40) calculated for daylight hours for each exposure ring are given in Table 1. Carbon dioxide was dispersed during daytime hours at 560 ppm, approximately 200 ppm above our ambient. The plants were established at 1 m × 1 m spacing across each ring to simulate an aggrading aspen forest. Details regarding the aspen clones are presented in Karnosky et al. (1998) and Sheng et al. (1997) and the planting and handling of the aspen clones is described in Karnosky et al. (1999).

The same plants were used for biomaterial measurements, determinations of visible leaf injury, antioxidant activities, chlorophyll, carotenoid, rubisco and starch concentrations, and gene expression levels (Rubisco small subunit and defense-related proteins), and therefore the data from different analyses are comparable. In addition, chlorophyll, carotenoid and rubisco concentrations were determined for the same leaf extraction samples.

## 2.2. Samples

Foliar samples from aspen were collected in late July of 1998 from each treatment ring as close as possible to mid-day (noon). Samples were collected from 6–10 trees in each treatment from each of the three replicates. The samples were immediately frozen in liquid nitrogen and stored at –80 °C until use. Leaf samples were collected from the developing leaf zone (LPI 3 to 5 (Larson and Isebrands, 1971)) and care was taken to collect same age leaf samples from each tree.

## 2.3. Sample extraction and analysis

Frozen leaves (0.25 g) were ground with liquid nitrogen and extracted briefly with 6 ml of cold 100 mM potassium phosphate buffer (pH 7.0) containing 0.1 mM EDTA and 1% (w/v) insoluble polyvinylpyrrolidone. The homogenate was centrifuged at 3000g (4 °C) for 2 min, 1 ml of supernatant was added to a 1 ml column of P2 Biogel (Bio Rad) and centrifuged at 600g for 1 min. Biogel purified samples were immediately assayed for SOD, APX and CAT activity. Samples assayed for GR activity were extracted as described above,

Table 1. Mean O<sub>3</sub> concentrations (ppb), total cumulative O<sub>3</sub> exposures over a threshold of 0 ppb (AOT00, ppm-h) and over a threshold of 40 ppb (AOT40, ppm-h) calculated for daylight hours during the 1998 growing season. Ambient O<sub>3</sub> was calculated for three points (North, East and South) within the exposure area

Month	O <sub>3</sub>	Ambient			Elevated O <sub>3</sub>			Elevated O <sub>3</sub> + CO <sub>2</sub>		
		North	East	South	1.3	2.3	3.3	1.4	2.4	3.4
May	Mean ppb	44	40	40	52	54	53	52	53	50
	AOT00 ppm-h	12.6	11.5	11.5	17.9	18.6	18.1	17.2	18.3	17.5
	AOT40 ppm-h	0.3	0.3	0.3	5.7	6.6	6.2	6.1	6.3	5.2
June	Mean ppb	35	31	32	55	57	55	55	57	55
	AOT00 ppm-h	12.5	11.3	11.7	19.7	20.5	19.7	19.9	20.3	19.7
	AOT40 ppm-h	0	0	0	6.7	7.9	6.9	7.3	7.9	6.7
July	Mean ppb	32	27	31	57	58	56	55	57	57
	AOT00 ppm-h	11.9	10	11.5	21.2	21.6	20.8	20.5	21.2	21.2
	AOT40 ppm-h	0	0	0	7.6	8.0	7.1	7.2	7.9	7.5
August	Mean ppb	44	37	41	59	58	59	64	62	57
	AOT00 ppm-h	13.6	11.4	12.6	18.4	18	18.3	19.8	19.1	17.8
	AOT40 ppm-h	0	0	0	7.0	8.1	7.0	9.2	8.4	6.5
May–August	Mean ppb	39	34	36	56	57	56	57	57	55
	AOT00 ppm-h	50.6	44.2	47.3	77.2	78.7	76.9	77.4	78.9	76.2
	AOT40 ppm-h	0.3	0.3	0.3	27.0	30.6	27.2	29.8	30.5	25.9

using 40 mM potassium phosphate buffer (pH 7.4) containing 1 mM EDTA, 3% (w/v) insoluble polyvinylpyrrolidone, and 40 mM 3-amino-1, 2,4-triazole. Homogenate was centrifuged at 6000g for 10 min (4 °C) and the supernatant was assayed for GR activity.

### 2.3.1. Antioxidant enzymes

Aspen clones (from samples collected in July) were analyzed to determine the levels of antioxidant enzymes. Biochemical analyses included enzyme assays for SOD (Dhindsa et al., 1981), ascorbate peroxidase (Chen and Asada, 1989), GR activity (Price et al., 1990), and catalase (Kato and Shimizu, 1987). All the biochemical analyses were normalized for age of the plants, weight of the sampled foliar material and for amount of total proteins or cell lysates used for each analysis. Assays were repeated on each extracted sample at least 3 times.

### 2.3.2. Rubisco, soluble protein, starch and pigments

For Rubisco and soluble protein assays, frozen leaf samples were weighed, and a crude extract was prepared using 2 ml of extraction buffer (Gezelius and Hallen, 1980). The amount of total Rubisco protein was determined by PAGE as described by Rintamäki et al. (1988), using purified Rubisco protein (Sigma Chemical Co.) as a standard. The areas and intensities of Rubisco bands were determined by scanning the gels with the Adobe PhotoShop Program (Version 5.0), and the Rubisco concentrations were calculated on a dry weight basis. An aliquot for soluble proteins was quantified as described by Pääkkönen et al. (1998). Chlorophyll analysis was done as per the protocol of Porra et al. (1989) and carotenoid measurements were done as per the protocol of Wellburn and Lichtenthaler (1984). In order to avoid variability between samples for Rubisco and protein analysis and chlorophyll and associated pigments, samples for pigments were taken from the crude extract before centrifugation and subjected to proper extraction protocol for either chlorophyll or carotenoid pigments. Leaf samples for starch determination were freeze-dried, milled, and analyzed by standard enzymatic techniques (Boehringer Kit for Food analysis).

### 2.3.3. Gene expression

Leaf material (1–2 g) was ground in liquid nitrogen, and added to the pre-warmed extraction buffer (100 mM Tris-HCl pH 8.0, 20 mM EDTA, 0.5 M NaCl, 2% PVP, 0.5% SDS, 0.5%  $\beta$ -mercaptoethanol). After three extractions with chloroform: IAA (24 : 1),  $1/5 \times$  volume of 10 M LiCl was added and the RNA was precipitated one hour on ice. The pellet was dissolved into DEPC-TE, extracted once more with chloroform:IAA, and reprecipitated in ethanol

at -80 °C overnight. The RNA pellet was purified with 70% ethanol, dried and resuspended in DEPC-treated water. RNA quality was checked by gel electrophoresis, and concentration determined by GeneQuant II RNA/DNA Calculator (Pharmacia Biotech). Single-stranded digoxigenin (DIG)-labeled DNA-probes were generated by PCR using DynaZyme DNA-polymerase (Finnzymes, Finland) and DIG-DNA labeling mixture (Boehringer Mannheim, Germany). Birch RbcS, Pal, Chs, and AccOx probes (see Pääkkönen et al., 1998) were used after careful optimization of hybridization conditions for aspen samples. RNA hybridizations were performed according to the DIG System User's Guide for Filter Hybridization (Boehringer Mannheim, Germany). Total RNA, 10 µg each per lane, was separated in formaldehyde containing 1% agarose gel. Equivalent loading of RNA was verified by SYBR-Green staining (Molecular Probes Inc., OR). After gel electrophoresis, RNA was transferred to nylon membranes overnight at room temperature with 10 × SSC (1.5 M NaCl, 150 mM sodium citrate pH 7.0). RNA was fixed to the filter by UV-cross-linking, followed by washing in 2 × SSC for 10 minutes. The filters were prehybridized for 1½ h at 48 °C in DIG Easy Hyb (Boehringer Mannheim, Germany) and hybridized overnight at 48 °C with denatured DIG-labeled purified DNA-probes. Filters were washed in 2 × SSC, 0.1% SDS at room temperature (2 × 5 min) and in 0.1 × SSC, 0.1% SDS at 68 °C (3 × 15 min). Hybridization signals were visualized using the DIG Luminescent Detection Kit (Boehringer Mannheim, Germany). Membranes were incubated in blocking reagent and treated with anti-DIG-alkaline phosphatase-conjugate. The chemiluminescent substrate was pipetted onto the filter and the light signal was recorded on X-ray films (Fig. 1). The relative intensities of stress protein inductions were determined by scanning the X-ray films and measuring the band densities with Adobe PhotoShop program (Version 5.0). The relative band densities were compared only within the film and the band intensities were indicated as percentage of the maximal intensity on the film.

#### 2.3.4. Growth analysis

Fifteen plants per clone per treatment were measured between 27 and 31 July 1998 for base diameter (mm), number of leaves per branch (3–6th branch from the top), mean leaf size (cm<sup>2</sup>; 9–11th leaf from the top) and proportion of visibly injured leaves (%; determined for the 3–6th branch from the top) as described (Karnosky and Steiner, 1980; Karnosky et al., 1996). Visible injuries were necrotic or dark-brown pigmented dots and flecks appearing on the upper leaf surface.




Figure 1. Effects of elevated O<sub>3</sub> and CO<sub>2</sub> alone or in combination on SOD, APX, GR and CAT activities in aspen clones 259, 216 and 271. Data for three replicate FACE rings/treatment were pooled and expressed as standard deviations.

Table 2. Effects of elevated O<sub>3</sub> and CO<sub>2</sub> alone and in combination on growth parameters in aspen (*Populus tremuloides*) clones. Significant block effects were not found, and therefore the data for three replicate FACE rings/treatment were pooled for further analyses. Anova, Tukey's test.  $P < 0.05$ ,  $n = 15$ . Values are means  $\pm$  S.E. Significant differences between the treatments are indicated by different letters

Response	Clone	Control	CO <sub>2</sub>	O <sub>3</sub>	CO <sub>2</sub> + O <sub>3</sub>
Base diameter, mm	259	18.8 $\pm$ 0.4 bc	20.3 $\pm$ 0.4 c	16.9 $\pm$ 0.5 a	20.1 $\pm$ 0.7 c
	216	22.2 $\pm$ 0.4 b	24.4 $\pm$ 0.4 c	18.9 $\pm$ 0.3 a	22.2 $\pm$ 0.4 b
	271	22.4 $\pm$ 0.5 b	22.6 $\pm$ 0.4 b	19.5 $\pm$ 0.5 a	21.9 $\pm$ 0.4 b
Number of leaves/branch	259	37.5 $\pm$ 1.0 b	36.5 $\pm$ 0.9 b	23.6 $\pm$ 0.6 a	23.2 $\pm$ 0.7 a
	216	34.8 $\pm$ 0.6 b	41.2 $\pm$ 1.2 c	23.9 $\pm$ 0.7 a	24.4 $\pm$ 0.7 a
	271	39.8 $\pm$ 0.9 b	39.9 $\pm$ 1.5 b	26.4 $\pm$ 0.7 a	25.5 $\pm$ 0.7 a
Mean leaf size, cm <sup>2</sup>	259	80.9 $\pm$ 4.7 a	67.8 $\pm$ 2.8 a	70.8 $\pm$ 5.3 a	70.1 $\pm$ 4.4 a
	216	67.4 $\pm$ 2.2 b	61.7 $\pm$ 3.4 ab	57.6 $\pm$ 2.1 ab	56.4 $\pm$ 2.9 a
	271	66.6 $\pm$ 3.0 b	63.1 $\pm$ 3.6 ab	59.8 $\pm$ 1.9 ab	54.3 $\pm$ 2.1 a
Injured leaves/branch, %	259	0.0 $\pm$ 0.0 a	0.0 $\pm$ 0.0 a	10.76 $\pm$ 0.65 b	8.11 $\pm$ 0.65 b
	216	0.0 $\pm$ 0.0 a	0.0 $\pm$ 0.0 a	8.42 $\pm$ 0.72 b	4.26 $\pm$ 0.54 b
	271	0.0 $\pm$ 0.0 a	0.0 $\pm$ 0.0 a	7.51 $\pm$ 0.71 b	3.30 $\pm$ 0.46 b

### 3. Results

#### 3.1. Leaf growth and visible injury

Ozone treated plants of all three clones had significantly smaller base diameters, average number of leaves per branch, and mean leaf sizes when compared to their respective controls, and similar results were obtained for average leaf sizes and leaves per branch of CO<sub>2</sub> + O<sub>3</sub> treated plants (Table 2). Average leaf sizes for clones 216 and 271 were most significantly affected by CO<sub>2</sub> + O<sub>3</sub>, indicating an increased sensitivity to O<sub>3</sub> in the presence of elevated CO<sub>2</sub> (Table 2). However, visible leaf injury (average number of injured leaves per branch as well as injury index per leaf measured as described by Karnosky and Steiner, 1980; Karnosky et al., 1996) was significantly higher for O<sub>3</sub> treated plants when compared to O<sub>3</sub> + CO<sub>2</sub> treated plants, and clone 259 showed the greatest amount of leaf injury (Table 2).

#### 3.2. Antioxidant activities

With the exception of samples from clone 259, SOD activities varied little, while APX, GR and CAT activities were generally highest for O<sub>3</sub> treated plants and lowest for CO<sub>2</sub> and O<sub>3</sub> + CO<sub>2</sub> treated plants for all three clones when compared to controls (Fig. 1).




Figure 2. Effects of elevated O<sub>3</sub> and CO<sub>2</sub> alone or in combination on chl a and b, total chl (mg/g dwt), carotenoid ( $\mu\text{g/g}$  dwt), total Rubisco (mg/g dwt), and starch (mg/g dwt) concentrations, and chl a/b ratio in aspen clones 259, 216 and 271. Anova, Tukey's test.  $P < 0.05$ ,  $n = 15$ . Values are means  $\pm$  S.E. Significant differences between the treatments are indicated by different letters. (Continued on next page).

### 3.3. Pigment, rubisco, and starch contents

Ozone, CO<sub>2</sub>, and O<sub>3</sub> + CO<sub>2</sub> treatments had significant effects on Chl a and Chl b content, whereas Rubisco content was not significantly affected (Fig. 2).




Figure 2. (Continued)

Chl a, chl b, and total chl (a + b) contents significantly decreased for all CO<sub>2</sub>, O<sub>3</sub>, and CO<sub>2</sub> + O<sub>3</sub> treated clones (Fig. 2). Chl a/b ratios only varied significantly for clone 259, with an increase (low chl b content) (Fig. 2). Total carotenoid and Rubisco concentrations generally decreased in CO<sub>2</sub>, O<sub>3</sub>, and CO<sub>2</sub> + O<sub>3</sub> treated plants. Starch concentrations for clones exposed to all three treatments, exhibited some trends, with CO<sub>2</sub>, O<sub>3</sub>, and CO<sub>2</sub> + O<sub>3</sub> plants generally having higher starch concentrations in clone 216 and lower concentrations in clones 259 and 271 when compared to controls (Fig. 2).

### 3.4. Transcription levels of *RbcS*, *Chs*, *Pal*, *AccOx*

Expression levels of *RbcS*, for all clones and all treatments, generally decreased compared to controls with the exception of O<sub>3</sub> treated clone 271 plants, which showed a significant increase (Figs. 3, 4). These results cor-




Figure 3. Transcript levels of Rubisco small subunit (RbcS), chalcone synthase (Chs), phenylalanine ammonia-lyase (Pal) and AccOx (means  $\pm$  S.E.) in leaves of aspen clones 259, 216 and 271. Maximum density of the band in each gel = 100. Kruskal-Wallis test. Significant differences between the treatments are indicated by different letters.

relate, most significantly for clone 259, with the Rubisco contents shown in Fig. 2, however, transcription levels do not appear to correlate with starch contents (Figs. 2, 3). Chs expression levels did not vary significantly, except for




Figure 4. Northern analysis of *RbcS*, *Pal*, *Chs*, and *AccOx* in aspen clones. Total RNA (10 µg each) from aspen clones 259, 216, and 271 was probed with single-stranded digoxigenin (DIG)-labeled Birch *RbcS*, *Pal*, *Chs*, and *AccOx* DNA probes.

O<sub>3</sub> treated clone 271 plants, which showed a significant increase compared to controls. Both clones 259 and 271 showed a significant decrease in total carotenoid contents under CO<sub>2</sub> and O<sub>3</sub> treatments (Fig. 2). The combination of CO<sub>2</sub> + O<sub>3</sub> had a rather dramatic effect in the decrease of carotenoid component on 259, but not on gene expression of *Chs*. *Pal* expression levels of all three clones were significantly increased by O<sub>3</sub> treatments and significantly decreased by CO<sub>2</sub> and CO<sub>2</sub> + O<sub>3</sub> treatments. Although the differences are not significant, changes in *AccOx* expression levels correlate with changes in *Pal* mRNA levels.

#### 4. Discussion

Polle et al. (1997) provided evidence that oxidative stress in beech leaves (*Fagus sylvatica* L.) increases when N and/or C assimilation was limited. Podila et al. (1998) found that clones grown in soils with the lowest average NH<sub>4</sub><sup>+</sup> and C:N ratios demonstrated a slight decrease in antioxidant activity. However, trends in the effects of elevated CO<sub>2</sub> on antioxidant activities could still

be seen in these aspen clones. Furthermore, Coleman et al. (1998) and Sober et al. (2001), found that N levels were linearly related to photosynthesis.

It has been reported that elevated CO<sub>2</sub> may result in decreased production of antioxidant metabolites (e.g., glutathione and ascorbate) and in antioxidant enzymes (Polle et al., 1993; Badiani et al., 1993; Schwanz et al., 1996; Polle et al., 1997; Polle and Pell, 1999) including aspen (Karnosky et al., 1998), while O<sub>3</sub> generally increases antioxidant activities. Furthermore, several reports have demonstrated that overexpression of SOD increases resistance to oxidative stress (Jansen et al., 1989; Sen Gupta et al., 1993a, 1993b), and that APX activities also increased in the leaves of transformed plants. In aspen (*Populus tremula* x *P. alba*), Tyystjärvi et al. (1999) found that an 5–8 fold increase in FeSOD expression was unable to protect the plants from photoinhibition, while tobacco plants, with a similar increase in GR expression levels, had a lower level of photoinhibition. Thus, the ascorbate-glutathione pathway may not be limited by the SOD catalyzed reduction of superoxide to H<sub>2</sub>O<sub>2</sub>, but instead by the conversion of H<sub>2</sub>O<sub>2</sub> to H<sub>2</sub>O by APX (and catalases) and the reduction of oxidized dehydroascorbate and glutathione by dehydroascorbate reductase and glutathione reductase. In our experiments we found that APX, CAT and GR activities for all three clones generally increased as a result of exposure to elevated O<sub>3</sub>, and decreased due to elevated CO<sub>2</sub> and CO<sub>2</sub> + O<sub>3</sub> levels, suggesting a suppression of the ascorbate-glutathione pathway by CO<sub>2</sub>.

A decrease in antioxidant activities would decrease the plants primary defense against AOS, and thus, an increase of oxyradicals within the chloroplast may account for the significant decreases observed for chl a and b and total carotenoid contents under elevated O<sub>3</sub>, especially since transcription levels of Chs were not significantly affected. This may also account for the decrease in Rubisco content. However, RbcS expression levels were also significantly lower in plants treated with CO<sub>2</sub>, O<sub>3</sub>, and CO<sub>2</sub> + O<sub>3</sub>. While it appears that more damage within the chloroplast occurs due to elevated CO<sub>2</sub> and O<sub>3</sub> levels, it was found in our work and in previous reports (Karnosky et al., 1999; Tjoelker et al., 1993; Volin et al., 1993) that CO<sub>2</sub> can reduce the amount of visible leaf injury. It has also been shown that CO<sub>2</sub> increases the light saturated photosynthetic rate for aspen (Karnosky et al., 1999; Volin and Reich, 1996; Volin et al., 1998); thus, it is possible that, in the presence of CO<sub>2</sub>, an increased rate of photosynthesis increases the concentration of available NADPH in the chloroplast, which can then be used by GR to reduce the glutathione pool and increase the flux through the ascorbate-glutathione cycle. However, leaf size was most significantly reduced for CO<sub>2</sub> + O<sub>3</sub> treated plants, suggesting that photosynthesis may be inhibited. Thus, it is also possible that an increase in AOS, due to suppression of antioxidants as discussed above, may be offset by a decrease in rate of photosynthesis which may only occur in CO<sub>2</sub> + O<sub>3</sub> treated plants, resulting in a decrease in both visible leaf injury and leaf growth.

Ozone sensitivity in *Populus* may be related to a lack of defense gene activation (Koch et al., 1998, 2000). Salicylic acid (SA) and jasmonic-acid-mediated (JA) signaling pathways, known mediators of pathogen and wound responses, may also be involved in responses to oxidative stress (Örvar et al., 1997; Koch et al., 1998, 2000; Rao et al., 1997, 2000). Our results indicate that Pal expression may be suppressed by CO<sub>2</sub> since Pal mRNA levels were lowest for CO<sub>2</sub> and CO<sub>2</sub> + O<sub>3</sub> treated plants (all clones). Suppression of the phenylpropanoid pathway, and a resulting decrease in various pathogen defense compounds, could increase the plants sensitivity to secondary pathogens especially when exposed to O<sub>3</sub> in combination with elevated CO<sub>2</sub>. This may explain the increased rate of infection by a sooty mold fungus, (*Alternaria* spp.) observed for all three clones grown at field sites with intermediate and high O<sub>3</sub> levels (Karnosky et al., 1999) and a 3- to 5-fold increase in *Melampsora* leaf rust in aspen exposed to O<sub>3</sub> or O<sub>3</sub> + CO<sub>2</sub> (Karnosky et al., 2001). Just as Karnosky et al. (1999) found clone 259 to be most dramatically affected by the secondary infection, we found that Pal expression was most significantly suppressed in clone 259. However, increased infection levels for clone 259 correlated well with increased visible leaf damage (Karnosky et al., 1999), suggesting that O<sub>3</sub>-induced foliar injury may result in a high susceptibility to secondary infections. It is also probable that clone 259 may be inherently less sensitive to SA-mediated signaling which may explain for its susceptibility to both O<sub>3</sub> and pathogens (Koch et al., 2000; Overmyer et al., 2000). On the other hand, clones 216 and 271 may be more sensitive to SA-mediated signaling resulting in launching a more effective defense against O<sub>3</sub>-mediated damage and/or pathogens.

The observation that AccOx levels were reduced under elevated CO<sub>2</sub> conditions for all three clones suggests that under elevated CO<sub>2</sub> conditions the production of ethylene may be reduced, leading to less visible injury due to oxidative damage as well as damage by pathogens. Recently, it has been shown that in *Arabidopsis* ethylene promotes SA-mediated (ozone-induced) lesion formation and the ozone-induced AccOx transcript levels match with ozone-induced ethylene production (Overmyer et al., 2000). Similarly, the reduction in AccOx levels under CO<sub>2</sub> + O<sub>3</sub> may have a role in reducing ethylene induced SA-mediated or O<sub>3</sub>-induced damage in clones 216 and 271. However, the low level of reduction for AccOx in clone 259 may yet lead to increased visible damage under CO<sub>2</sub> + O<sub>3</sub> conditions.

## 5. Conclusions

Although responses to elevated CO<sub>2</sub>, O<sub>3</sub>, and CO<sub>2</sub> + O<sub>3</sub> were similar in all clones, clone 259 showed the highest sensitivity to all treatments. Our results

suggest that elevated CO<sub>2</sub> levels may exacerbate the harmful effects of O<sub>3</sub> by suppressing the ascorbate-glutathione and Pal pathways, resulting in increased cellular damage as a result of increasing AOS levels and a reduced resistance to secondary infections. Degradation due to an increase in AOS, resulting from a decrease in antioxidant activities, was most likely responsible for the observed decrease in chlorophyll and carotenoid contents, while decreased expression levels of RbcS correlated well with decreased Rubisco contents.

### Acknowledgements

This research was partially supported by the USDA Forest Service Global Change Program, the US Department of Energy, Office of Biological and Environmental Research (BER) (DE-FG02-95ER62125 and DE-FG02-93ER6166), and the Brookhaven National Labs/US Department of Energy (725079), the National Science Foundation (DBI-9601942, DBI-9601942, and IBN-9652675), the National Council of Air and Stream Improvement of the Pulp and Paper Industry (NCASI), and the Academy of Finland.

### References

- Badiani, M., D'Annibale, A., Paolacci, A.R., Miglietta, F., Raschi, A., 1993. The antioxidant status of soybean (*Glycine max*) leaves grown under natural CO<sub>2</sub> enrichment in the field. *Aust. J. Plant Physiol.* 20, 275–284.
- Barnola, J.M., Anklin, M., Porheron, J., Raynaud, D., Schwander, J., Stauffer, B.T.I., 1995. CO<sub>2</sub> evolution during the last millennium as recorded by Antarctic and Greenland ice. *Tellus B* 47, 264–272.
- Broadmeadow, M.S.J., Heath, J., Randle, T.J., 1999. Environment limitations to O<sub>3</sub> uptake—some key results from young trees growing at elevated CO<sub>2</sub> concentrations. In: Sheppard, L.J., Cape, J.N. (Eds.), *Forest Growth Responses to the Pollution Climate of the 21st Century*. Water Air Soil Pollut., pp. 299–310.
- Chameides, W.L., Kasibhatla, P.S., Yienger, J., Levy II, H., 1995. Growth of continental-scale metro-agro-plexes, regional ozone pollution, and world food production. *Science* 264, 74–77.
- Chen, G.S., Asada, K., 1989. Ascorbate peroxidase in the tea leaves: Occurrence of two isozymes and the differences in their enzymatic and molecular properties. *Plant Cell Physiol.* 30, 987–998.
- Coleman, M.D., Dickson, R.E., Isebrands, J.G., 1998. Growth and physiology of aspen supplied with different fertilizer addition rates. *Physiol. Plant.* 103, 513–526.
- Dickson, R.E., Coleman, M.D., Riemenschneider, D.E., Isebrands, J.G., Hogan, G.D., Karnosky, D.F., 1998. Growth of five hybrid poplar genotypes exposed to interacting elevated CO<sub>2</sub> and O<sub>3</sub>. *Can. J. Forest Res.* 28, 1706–1716.
- Dickson, R.E., Lewin, K.F., Isebrands, J.G., Coleman, M.D., Heilman, W.E., Riemenschneider, D.E., Sober, J., Host, G.E., Zak, D.F., Hendrey, G.R., Pregitzer, K.S., Karnosky, D.F., 2000. Forest atmosphere carbon transfer storage-II (FACTS II)—The aspen free-air CO<sub>2</sub> and O<sub>3</sub> enrichment (FACE) project in an overview. USDA Forest Service North Central Research Station. General Tech. Rep. NC-214.

- Dhindsa, R.S., Plumb-Dhindsa, P., Thorpe, T.A., 1981. Leaf senescence correlated with increased levels of membrane permeability and lipid peroxidation, and decreased levels of superoxide dismutase and catalase. *J. Exp. Bot.* 32, 93–101.
- Gezelius, K., Hallen, M., 1980. Seasonal variation in ribulose biphosphate carboxylase activity in *Pinus sylvestris*. *Physiol. Plant.* 48, 88–98.
- Grams, T.E.E., Anegg, S., Haberle, K.H., Langebartels, C., Maytessek, R., 1999. Interactions of chronic exposure to elevated CO<sub>2</sub> and O<sub>3</sub> levels in the photosynthetic light and dark reactions of European beech (*Fagus sylvatica*). *New Phytol.* 144, 95–107.
- Jansen, A.K.J., Shaaltiel, Y., Kazzes, D., Canaani, O., Malkin, S., Gessel, J., 1989. Increased tolerance to photoinhibitory light in paraquat-resistant *Conyza bonariensis* measured by photoacoustic spectroscopy and <sup>14</sup>CO<sub>2</sub>-fixation. *Plant Physiol.* 91, 1174–1178.
- Karnosky, D.F., Steiner, K.C., 1980. Provenance and family variation in response of *Fraxinus americana* and *F. pennsylvanica* in response to ozone and sulfur dioxide. *Phytopathology* 71, 804–807.
- Karnosky, D.F., Gagnon, Z.E., Dickson, R.E., Coleman, M.D., Lee, H., Isebrands, J.G., 1996. Changes in growth, leaf abscission and biomass associated with seasonal tropospheric ozone exposures of *Populus tremuloides* clones and seedlings. *Can. J. Forest Res.* 26, 23–37.
- Karnosky, D.F., Podila, G.K., Gagnon, Z., Pechter, P., Akkapeddi, A., Sheng, Y., Riemenschneider, D.E., Coleman, M., Dickson, R.E., Isebrands, J.G., 1998. Genetic control of responses to interacting tropospheric ozone and CO<sub>2</sub> in *Populus tremuloides*. *Chemosphere* 36, 807–812.
- Karnosky, D.F., Mankovska, B., Percy, K., Dickson, R.E., Podila, G.K., Hendrey, G., Coleman, M.D., Kubiske, M., Pregitzer, K.S., Sober, J., Isebrands, J.G., 1999. Effects of tropospheric O<sub>3</sub> and interaction with CO<sub>2</sub>: Comparison of open-top chamber, field and FACE results with trembling aspen genotypes differing in O<sub>3</sub> sensitivity. *J. Air Water Soil Pollut.* 116, 311–322.
- Karnosky, D.F., Percy, K.E., Xiang, B., Callan, B., Noormets, A., Mankovska, B., Hopkin, A., Sober, J., Jones, W., Dickson, R.E., Isebrands, J.G., 2001. Interacting CO<sub>2</sub>-tropospheric O<sub>3</sub> and predisposition of aspen (*Populus tremuloides* Michx.) to infection by *Melampsora medusae* rust. *Global Change Biol.* 8, 329–338.
- Kato, M., Shimizu, S., 1987. Chlorophyll metabolism in higher plants VII. Chlorophyll degradation in senescing tobacco leaves phenolic-dependent peroxidative degradation. *Can. J. Bot.* 65, 729–735.
- Kellomaki, S., Wang, K.Y., 1997. Effects of elevated O<sub>3</sub> and CO<sub>2</sub> concentrations on photosynthesis and stomatal conductance in Scots pine. *Plant Cell Environ.* 20, 995–1006.
- Koch, R.K., Scherzer, A.J., Eshita, S.M., Davis, K.R., 1998. Ozone sensitivity in hybrid poplar is correlated with a lack of defense-gene activation. *Plant Physiol.* 118, 1243–1252.
- Koch, J.R., Creelman, R.A., Eshita, S.M., Seskar, M., Mullet, J.E., Davis, K.R., 2000. Ozone sensitivity in hybrid poplar correlates with insensitivity to both salicylic acid and jasmonic acid. The role of programmed cell death in lesion formation. *Plant Physiol.* 123, 487–496.
- Kull, O., Sober, A., Coleman, M.D., Dickson, R.E., Isebrands, J.G., Gagnon, Z., Karnosky, D.F., 1996. Photosynthetic response of aspen clones to simultaneous exposures of ozone and CO<sub>2</sub>. *Can. J. Forest Res.* 16, 639–648.
- Larson, P.R., Isebrands, J.G., 1971. The plastochron index as applied to developmental studies of cottonwood. *Can. J. Forest Res.* 1, 1–11.
- Matyssek, R., Innes, J.L., 1999. Ozone—a risk factor for trees and forests in Europe? *J. Water Air Soil Pollut.* 116, 199–226.
- McKee, I.F., Farage, P.K., Long, S.P., 1995. The interactive effects of elevated CO<sub>2</sub> and O<sub>3</sub> concentration on photosynthesis in spring wheat. *Photosynth. Res.* 45, 111–119.
- Mortensen, L.M., 1995. Effect of carbon dioxide concentration on biomass production and partitioning in *Betula pubescens* Ehrh. Seedlings at different ozone and temperature regimes. *Environ. Pollut.* 87, 337–343.

- Niewiadomska, E., Gaucher-Velleux, C., Chevrier, N., Maufette, Y., Dizengremel, P., 1999. Elevated CO<sub>2</sub> does not provide protection against ozone considering activity of several antioxidant enzymes in the leaves of sugar maple. *J. Plant Physiol.* 155, 70–77.
- Noormets, A., Podila, G.K., Karnosky, D.F., 2000. Rapid response of antioxidant enzymes to O<sub>3</sub>-induced oxidative stress in *Populus tremuloides* clones varying in O<sub>3</sub> tolerance. *Forest Genetics* 7, 339–342.
- Örvar, B.J., McPherson, J., Ellis, B.E., 1997. Pre-activating wounding response in tobacco prior to high-level ozone exposure prevents necrotic injury. *Plant J.* 11, 203–212.
- Overmyer, K., Tuominen, H., Kettunen, R., Betzb, C., Langebartels, C., Sandermann Jr., H., Kangasjärvi, J., 2000. Ozone-sensitive arabidopsis *rcd1* mutant reveals opposite roles for ethylene and jasmonate signaling pathways in regulating superoxide-dependent cell death. *Plant Cell* 12, 1849–1862.
- Pääkkönen, E., Vahala, J., Pohjola, M., Holopainen, T., Kärenlampi, L., 1998. Physiological, stomatal and ultrastructural ozone responses in birch (*Betula pendula* Roth.) are modified by water stress. *Plant Cell Environ.* 21, 671–684.
- Podila, G.K., Wustman, B.A., Wang, Y., Zak, D.R., Sober, J., Hendrey, G.R., Isebrands, J.G., Pregitzer, K., Karnosky, D.F., 1998. Effects of elevated CO<sub>2</sub> on the antioxidant potential of aspen clones varying in O<sub>3</sub> sensitivity: Results from a free air CO<sub>2</sub> exposure (FACE) project. In: Proc. 18th International Meeting for Specialists in Air Pollution Effects on Forest Ecosystems, Edinburgh, Scotland (Abstract, p. 97).
- Polle, A., Pfirrmann, T., Chakrabarti, S., Rennenberg, H., 1993. The effects of enhanced ozone and enhanced carbon dioxide concentrations on biomass, pigments and antioxidative enzymes in spruce needles (*Picea abies* L.). *Plant Cell Environ.* 16, 311–316.
- Polle, A., Eiblmeier, M., Sheppard, L., Murray, M., 1997. Responses of antioxidant enzymes to elevated CO<sub>2</sub> in leaves of beech (*Fagus sylvatica* L.) seedlings grown under a range of nutrient regimes. *Plant Cell Environ.* 20, 1317–1321.
- Polle, A., Pell, E., 1999. Role of carbon dioxide in modifying the plant response to ozone. In: Luo, Y., Mooney, H.A. (Eds.), *Carbon Dioxide and Environmental Stress*. Academic Press, New York, pp. 193–213.
- Porra, R.J., Thompson, W.A., Kriedemann, P.E., 1989. Determination of accurate extinction coefficients and simultaneous equations for assaying chlorophylls a and b extracted with four different solvents: verification of the concentration of chlorophyll standards by atomic absorption spectroscopy. *Biochem. Biophys. Acta* 975, 384–394.
- Price, A., Lucas, P.W., Lea, P.J., 1990. Age-dependent damage and glutathione metabolism in ozone-fumigated barley: A leaf section approach. *J. Exp. Bot.* 41, 1309–1317.
- Rao, M.V., Hale, B.A., Ormrod, D.P., 1995. Amelioration of ozone-induced oxidative damage in wheat plants grown under high carbon dioxide. *Plant Physiol.* 109, 421–432.
- Rao, M.V., Lee, H., Creelman, R.A., Mullet, J.E., Davis, K.R., 2000. Jasmonic acid signaling modulates ozone-induced hypersensitive cell death. *Plant Cell* 12, 1633–1646.
- Rao, M.V., Paliyath, G., Ormrod, D.P., Murr, D.P., Watkins, C.B., 1997. Influence of salicylic acid on H<sub>2</sub>O<sub>2</sub> production, oxidative stress, and H<sub>2</sub>O<sub>2</sub>-metabolizing enzymes. *Plant Physiol.* 115, 137–149.
- Reilly, J., Prinn, R., Harnisch, J., Fitzmaurice, J., Jacoby, H., Kicklighter, D., Mellilo, J., Stone, P., Sokolov, A., Wang, C., 1999. Multi-gas assessment of the Kyoto Protocol. *Nature* 401, 549–555.
- Rintamäki, E., Keys, A.J., Parry, M.A.J., 1988. Comparison of the specific activity of ribulose-1,5-bisphosphate carboxylase-oxygenase from some C<sub>3</sub> and C<sub>4</sub> plants. *Physiol. Plant.* 74, 326–331.
- Schwanz, P., Kimball, B.A., Idso, S.B., Hendrix, D.L., Polle, A., 1996. Antioxidants in sun and shade leaves of sour orange trees (*Citrus aurantium*) after long-term acclimation to elevated CO<sub>2</sub>. *J. Exp. Bot.* 47, 1941–1950.

- Sehmer, L., Fontaine, V., Antoni, F., Dizengremel, P., 1998. Effects of ozone and elevated atmospheric carbon dioxide on carbohydrate metabolism of spruce needles. Catabolic and detoxification pathways. *Physiol. Plant.* 102, 605–611.
- Sen Gupta, A., Heinen, J.L., Holaday, A.S., Burke, J.J., Allen, R.D., 1993a. Increased resistance to oxidative stress in transgenic plants that overexpress chloroplastic Cu/Zn superoxide dismutase. *Proc. Nat. Acad. Sci. USA* 90, 1629–1633.
- Sen Gupta, A., Webb, R.P., Holaday, A.S., Allen, R.D., 1993b. Overexpression of superoxide dismutase protects plants from oxidative stress: Induction of ascorbate peroxidase in superoxide dismutase-overexpressing plants. *Plant Physiol.* 103, 1067–1073.
- Sheng, Y., Podila, G.K., Karnosky, D.F., 1997. Differences in O<sub>3</sub>-induced SOD and glutathione antioxidant expression in tolerant and sensitive aspen (*Populus tremuloides* Michx.) clones. *Forest Genetics* 4, 25–33.
- Sober, A., Noormets, A., Kull, O., Isebrands, J.G., Dickson, R.E., Sober, J., Karnosky, D.F., 2001. Photosynthetic parameters in aspen grown with interacting elevated CO<sub>2</sub> and tropospheric ozone concentrations as affected by leaf nitrogen. *Tree Physiol.* (in press).
- Tjoelker, M.G., Volin, J.C., Oleksyn, J., Reich, P., 1993. Light environment alters response to ozone stress seedlings of *Acer saccharum* Marsh. and hybrid *Populus* L. I. *In situ* net photosynthesis, dark respiration and growth. *New Phytol.* 124, 627–636.
- Tyystjärvi, E., Riikonen, M., Arisi, A.-C.M., Kettunen, R., Jouanin, L., Foyer, C.H., 1999. Photoinhibition of photosystem II in tobacco plants overexpressing glutathione reductase and poplars overexpressing superoxide dismutase. *Physiol. Plant.* 105, 409–416.
- Utrainen, J., Janhunen, S., Helmisaari, H.S., Holopainen, T., 2000. Biomass allocation, needle structural characteristics and nutrient composition in Scots pine seedlings exposed to elevated CO<sub>2</sub> and O<sub>3</sub> concentrations. *Trees* 14, 475–484.
- Volin, J.C., Tjoelker, M.G., Oleksyn, J., Reich, P.B., 1993. Light environment alters response to ozone stress in seedlings of *Acer saccharum* Marsh. and hybrid *Populus* L. II. Diagnostic gas exchange and leaf chemistry. *New Phytol.* 124, 637–646.
- Volin, J.C., Reich, P.B., 1996. Interaction of elevated CO<sub>2</sub> and O<sub>3</sub> on growth, photosynthesis and respiration of three perennial species grown in low and high nitrogen. *Physiol. Plant.* 97, 674–684.
- Volin, J.C., Reich, P.B., Givnish, T.J., 1998. Elevated carbon dioxide ameliorates the effects of ozone on photosynthesis and growth: species respond similarly regardless of photosynthetic pathway or plant functional group. *New Phytol.* 138, 315–325.
- Wellburn, A.R., Lichtenthaler, H., 1984. Formulae and program to determine total carotenoids and chlorophylls a and b of leaf extracts in different solvents. In: Sybesma, C. (Ed.), *Adv. Photosynth. Res.*, Vol. 2, pp. 9–12.