

## List of Figures

Fig. 2-1:	Flowchart of the Impact Pathway Approach including monetary valuation .....	13
Fig. 2-2:	Maximal time scales between contamination of different media leading to exposures via inhalation and/or ingestion and impacts on human health (cliparts by Corel Corporation, 1999, 2002).....	16
Fig. 2-3:	Options for the combination of the spatial scope, lateral spatial resolution and compartmentalisation of an environmental fate (and exposure) model (clipart by Corel Corporation, 2002) .....	25
Fig. 4-1:	Conceptual structure of the environmental fate and exposure assessment of the WATSON model and its linkage to the air quality model (arrows connecting boxes denote a substance's environmental pathway, arrows not connecting boxes indicate ultimate removal processes from the model's scope) .....	66
Fig. 4-2:	Spatial resolution of the WATSON model based on watersheds; data taken from EROS Data Center (1996) and adjusted (see text).....	80
Fig. 4-3:	Spatial resolution of the WATSON model based on watersheds which are further subdivided in the case of larger catchments; data taken from EROS Data Center (1996) and adjusted (see text) .....	81
Fig. 4-4:	Organisation of the Rhine catchment including the Meuse river according to the Pfafstetter code (note the Rhine catchment is identified by "914" at the continental scale, the shown digits constitute the fourth level subdivision, i.e., "914x"; the general Pfafstetter coding principle is also shown at the top).....	82
Fig. 5-1:	Distribution of the predominance of arable land (left) and pastures/grasslands (right) in the different zones distinguished by WATSON .....	108
Fig. 5-2:	Distribution of the predominance of (semi-) natural ecosystems (left) and non-vegetated land (right) in the different zones distinguished by WATSON .....	109

- Fig. 5-3: Distribution of the predominance of impervious surfaces (left) and glaciers (right) in the different zones distinguished by WATSON ..... 110
- Fig. 6-1: Distribution of the predominance of freshwater bodies in the different zones distinguished by WATSON (note that the Black Sea and the Caspian Sea are presently not modelled)..... 137
- Fig. 6-2: Lake Vänern in southern Sweden as an example of a sub-division of larger lakes according to the spatial differentiation as provided by HYDRO1k basin dataset (EROS Data Center, 1996; dark grey: lakes; light grey: the Götälv catchment; water grossly flowing from north-east to south-west; lakes fully contained in one zone are also shown) ..... 150
- Fig. 8-1: Temporal structure of the distribution of 6.09 and 15.95 years of life lost (YOLL) due to an exposure towards a pollutant at time 0 years without a minimum latency time at the population level ..... 198
- Fig. 8-2: Temporal structure of the distribution of 15.95 years of life lost (YOLL) due to an exposure towards a pollutant at time 0 years according to three different assumptions with respect to latency at the population level: no minimum latency, minimum latency of 10 years and absolute latency of 10 years (black, grey and white circles, respectively)..... 198
- Fig. 9-1: Comparison of atmospheric depositions in micrograms per square metre and year based on air quality modelling by the Windrose Trajectory Model for emissions in year 1998 (left) and moss concentrations in micrograms per gram for the years 2000/2001 (right, taken from Buse et al. (2003) and including data for Iceland, with permission) for arsenic (top) and cadmium (bottom)..... 234
- Fig. 9-2: Comparison of atmospheric depositions in micrograms per square metre and year based on air quality modelling by the Windrose Trajectory Model for emissions in year 1998 (left) and moss concentrations in micrograms per gram for the years 2000/2001 (right, taken from Buse et al. (2003) and including data for Iceland, with permission) for chromium ..... 235
- Fig. 9-3: Comparison of arsenic minimum, median and maximum concentrations as predicted by the environmental fate and exposure sub-models with reported concentrations in environmental media (top) and food-stuff (bottom, cf. Table C-9); model estimates based on the 'food removal' scenario described in Table 9-1 resulting from a 100 year continuous release according to the pan-European emission scenario for 1990 (cf. sections 11.1 and 11.2; different units; note the logarithmic scale, horizontal bars indicate reported detection limits) ..... 238

- Fig. 9-4: Comparison of cadmium minimum, median and maximum concentrations as predicted by the environmental fate and exposure models with reported concentrations in environmental media (top) and foodstuff (bottom, cf. Table C-10); model estimates based on the 'food removal' scenario described in Table 9-1 resulting from a 100 year continuous release according to the pan-European emission scenario for 1990 (cf. sections 11.1 and 11.2; different units; note the logarithmic scale, horizontal bars indicate reported detection limits)..... 239
- Fig. 9-5: Comparison of chromium minimum, median and maximum concentrations as predicted by the environmental fate and exposure models with reported concentrations in environmental media (top) and foodstuff (bottom, cf. Table C-11); model estimates based on the 'food removal' scenario described in Table 9-1 resulting from a 100 year continuous release according to the pan-European emission scenario for 1990 (cf. sections 11.1 and 11.2; different units; note the logarithmic scale, horizontal bars indicate reported detection limits) ..... 240
- Fig. 9-6: Comparison of lead minimum, median and maximum concentrations as predicted by the environmental fate and exposure models with reported concentrations in environmental media (top) and foodstuff (bottom, cf. Table C-12); model estimates based on the 'food removal' scenario described in Table 9-1 resulting from a 100 year continuous release according to the pan-European emission scenario for 1990 (cf. sections 11.1 and 11.2; different units; note the logarithmic scale, horizontal bars indicate reported detection limits; predicted beef concentrations are compared to a measured value for pork) ..... 241
- Fig. 9-7: Effective Intake Fraction for cadmium due to the ingestion of food according to the sensitivity scenarios after 25 years, 100 years and at steady state (top) and the development within the first 500 years after the pulse emission (bottom); pan-European emissions to air in 1990 ..... 250
- Fig. 9-8: Concentration distribution of cadmium in agricultural soil at steady-state due to 1990 pan-European emissions to air according to the 'low resolution' (top) and 'simple high resolution' (bottom) scenarios [mg/kg] ..... 252
- Fig. 9-9: Concentration distribution of cadmium in freshwater at steady-state due to 1990 pan-European emissions to air according to the 'low resolution' (top) and 'simple high resolution' (bottom) scenarios [mg/l] ..... 253
- Fig. 9-10: Factors obtained by relating the concentrations of cadmium according to the 'low resolution' scenario to those assessed by the 'simple high

- resolution' scenario at steady-state in the freshwater (left) and agricultural soil compartments (right) [-] (pan-European emissions to air in 1990) ..... 254
- Fig. 9-11: Factors by which the freshwater concentration of cadmium according to the 'rivers from lakes distinguished' scenario deviates from those assessed by the 'simple high resolution' scenario at steady-state [-] (pan-European emissions to air in 1990) ..... 255
- Fig. 9-12: Atmospheric deposition of cadmium according to the pan-European emission scenario for 1990 [ $\mu\text{g}/\text{m}^2/\text{yr}$ ] ..... 256
- Fig. 9-13: Effective Intake Fraction for chromium due to ingestion of food according to the sensitivity analysis after 25 years, 100 years and at steady state (top) and the development within the first 500 years after the pulse emission (bottom); pan-European emissions to air in 1990 ..... 264
- Fig. 9-14: Relative contribution of the different food items to the effective Intake Fraction (ingestion) of chromium after 25 years (top) and time-integrated (bottom) for a one year pulse emission according to the pan-European emission scenario to air in 1990 (cliparts by Corel Corporation, 2002) ..... 265
- Fig. 10-1: Effective Intake Fraction of selected trace elements via inhalation after one year and via ingestion of food after 10 and 100 years, and time-integrated for a one year pulse emission from the Belgian ('BE', top) and French site ('FR', bottom); note the logarithmic scale [ $\text{kg}_{\text{intake}}$  per  $\text{kg}_{\text{released}}$ ] ..... 288
- Fig. 10-2: Effective Intake Fraction of selected trace elements via inhalation after one year and via ingestion of food after 10 and 100 years, and time-integrated for a one year pulse emission from the German ('DE', top) and UK site ('UK', bottom); note the logarithmic scale [ $\text{kg}_{\text{intake}}$  per  $\text{kg}_{\text{released}}$ ] ..... 289
- Fig. 10-3: Relative contribution of the different food items to the effective Intake Fraction (ingestion) of selected trace elements after 10 years and time-integrated for a one year pulse emission from the Belgian ('BE', top) and French site ('FR', bottom, cliparts by Corel Corporation, 2002) ..... 290
- Fig. 10-4: Relative contribution of the different food items to the effective Intake Fraction (ingestion) of selected trace elements after 10 years and time-integrated for a one year pulse emission from the German ('DE', top) and the UK site ('UK', bottom, cliparts by Corel Corporation, 2002) ..... 291
- Fig. 11-1: Cadmium concentrations in arable land after 10 years (top left), 100 years (top right), 1000 years (bottom left) and at steady-state (bottom

	right) according to the pan-European emission scenario for 1990 (continuous releases) [mg/kg].....	303
Fig. 11-2:	Arsenic concentrations in arable land after 10 years (top left), 100 years (top right), 1000 years (bottom left) and at steady-state (bottom right) according to the pan-European emission scenario for 1990 (continuous releases) [mg/kg].....	304
Fig. 11-3:	Arsenic concentrations in freshwater bodies after 10 years (top left), 100 years (top right), 1000 years (bottom left) and at steady-state (bottom right) according to the pan-European emission scenario for 1990 (continuous releases) [pg/l].....	305
Fig. 11-4:	Development of arsenic concentrations towards the steady-state in the Hron River catchment in central Slovakia according to the pan-European emission scenario for 1990 (continuous releases); the values are given relative to the steady-state situation (cf. Table 11-1) [-].....	306
Fig. 11-5:	Effective Intake Fraction of selected trace elements via inhalation after one year and via ingestion of food after 10 and 100 years, and time-integrated for a one year pulse emission according to the pan-European emission scenario in 1990 (note the logarithmic scale) [ $\text{kg}_{\text{intake}}$ per $\text{kg}_{\text{released}}$ ] .....	311
Fig. 11-6:	Relative contribution of the different food items to the effective Intake Fraction (ingestion) of selected trace elements after 10 years (top) and time-integrated (bottom) for a one year pulse emission according to the pan-European emission scenario to air in 1990 (cliparts by Corel Corporation, 2002).....	312
Fig. A-1:	Masses with respect to removal due to harvest resulting in the environmental fate and exposure model in the dynamic case.....	439
Fig. B-1:	Area for which concentrations and depositions are calculated on the EMEP 50 km grid within the single and multi source EcoSense Europe version (European Commission, 1999a; Friedrich and Bickel, 2001a) .....	455
Fig. B-2:	Properties of top soil in terms of soil reaction (pH, left) and organic carbon reservoir (right, [ $\text{kg}_{\text{carbon}}/\text{m}^2$ ]; taken from Batjes, 1996) ...	479
Fig. B-3:	Long-term values for runoff from land (top) and ground water recharge (bottom) in the area of interest according to Döll and co-workers (Döll and Lehner, 2002; Döll et al., 2002, 2003) [mm/yr] .....	486
Fig. B-4:	Subdivision of the geographical scope of the model into administrative units. Countries are distinguished by different shades of grey (see Fig. B-1 for the model's boundaries).....	511
Fig. B-5:	Example on the deviation of the food supply data (Food and Agriculture Organization of the United Nations - Statistics Division, 2002a)	

from the consumption data by Euromonitor (1992): EU15 countries in  
1990 ..... 518