

List of Tables

Table 2-1:	Summary of the pollutants currently considered in different EcoSense Europe versions	14
Table 2-2:	Air quality models implemented in EcoSense.....	15
Table 2-3:	Attempt to structure the implications of different substance properties, reaction chemistry and modes-of-entry on model design	20
Table 3-1:	Characteristics of global multi-zonal multimedia environmental fate models without exposure assessment	35
Table 3-2:	Characteristics of a gridded multi-zonal multimedia environmental fate model for Europe (without exposure assessment; Prevedouros et al., 2004).....	36
Table 3-3:	Characteristics of multi-zonal multimedia environmental fate models applicable to particular regions of the world.....	38
Table 3-4:	Characteristics of IMPACT 2002, a multi-zonal multimedia environmental fate and exposure model.....	41
Table 3-5:	Characteristics of oligo-zonal multimedia environmental fate and exposure models	43
Table 3-6:	Characteristics of multimedia exposure approaches for trace elements/radionuclides	52
Table 4-1:	Treatment of different particle size classes by the Windrose Trajectory Model (WTM).....	67
Table 4-2:	Feedback fractions of selected substances (Margni, 2002)	69
Table 4-3:	Process formulations determining the (exogenous) inputs into the water and soil compartments	72
Table 5-1:	Terrestrial compartments distinguished according to qualitative criteria; their area shares in the geographical scope of the model are also given (derived from data presented in section B.3)	90
Table 5-2:	Overview on different soil depths adopted by selected multimedia models.....	92
Table 5-3:	Soil characteristics according to different multimedia models	95

Table 5-4:	Overview on different soil-related processes considered by selected multimedia models	97
Table 5-5:	Process formulations for terrestrial compartments as used in the present assessment	98
Table 5-6:	Relative erodibility of different land covers according to Golubev (1982) and their assignment to compartments as used in this study	102
Table 5-7:	Compartment-specific water soil erosion weights and velocities depending on the number of soil compartments distinguished which show the related process	103
Table 5-8:	Non-exhaustive overview on existing plant models in the field of multimedia models	114
Table 5-9:	Process formulations for terrestrial plants of agricultural use as used in the present assessment	132
Table 6-1:	Characteristics of solids in the freshwater environment as used in the presented methodology	140
Table 6-2:	Non-exhaustive overview about processes considered for the freshwater compartment by various multimedia models (note: chemical transformations are not listed)	142
Table 6-3:	Process formulations for the aquatic environment as used in the present assessment	143
Table 6-4:	Particle mass balance for surface freshwater assumed in this study differentiated into a pure river and a pure lake situation	147
Table 7-1:	Parameter values adopted in the exposure assessment deviating from those recommended by the United States - Environmental Protection Agency (1998) for ingestion	157
Table 7-2:	Exposure pathway formulations for ingestion exposures as used in the exposure assessment	158
Table 7-3:	Estimation of ED_{10} from threshold effect measures	170
Table 7-4:	Estimation of the β_{ED10} slope factor based on linear exposure-response information (cf. footnote 18)	171
Table 7-5:	International Life Sciences Institute classification scheme for human health impact categories (Burke et al., 1996 taken from Owens, 2001 and Pennington et al., 2002)	177
Table 7-6:	Cancer effect-related β_{ED10} slope factors and physical impacts for mortality (YOLL) and morbidity (YLD) due to inhalation and ingestion exposure of selected trace elements	178
Table 7-7:	Non-cancer effect-related β_{ED10} slope factors and aggregated physical impacts for mortality and morbidity in terms of DALYs due to inhalation and ingestion exposure of selected trace elements	180

Table 7-8:	Definitions of disability weighting in the Global Burden of Disease Study according to Murray (1994)	184
Table 8-1:	Declining discount rate scheme suggested by Weitzmann (1999)	190
Table 8-2:	Monetary values used for the valuation of the costs of illness (COI) for the endpoints considered in this study	196
Table 8-3:	Monetary values per YOLL when discounting at a rate of 3 % according to the approach followed in this study [€ ₂₀₀₀ per YOLL]	199
Table 8-4:	Implications of the pain and suffering-related monetary value for cancers according to the DALY and the ExternE approach for a discount rate of 0 %.....	200
Table 8-5:	Impact of choosing the DALY or the ExternE approach with respect to valuing pain and suffering-related monetary valuation when discounting at a rate of 0 %.....	201
Table 8-6:	Monetary values per incidence of a disease by valuation approach and discount rate as used in the present study [€ ₂₀₀₀ per case]...	203
Table 9-1:	Scenarios evaluated with respect to the spatial resolution, the compartments distinguished and adapted processes.....	244
Table 9-2:	Contribution of the different food items to the Intake Fraction (last row) of cadmium for time-integrated ingestion exposures according to the sensitivity scenarios (pan-European emissions to air in 1990)	248
Table 9-3:	Contribution of the different food items to the Intake Fraction (last row) of cadmium for ingestion exposures after 25 years according to the sensitivity scenarios (pan-European emissions to air in 1990, only taking place in the first year)	249
Table 9-4:	Maximum concentrations in agricultural produce at steady-state for air emissions in 1990 according to the exposure assessments as given by International Atomic Energy Agency (2001) and United States - Environmental Protection Agency (1998) for Europe	259
Table 9-5:	Assigned pH values in the 'compartmental pH variation' sensitivity case	261
Table 9-6:	Components of the effective Intake Fraction for chromium due to the ingestion of different food items according to the considered sensitivity cases after 25 years and time-integrated for the pan-European emission scenario to air in 1990 (emissions only take place in the first year) [kg _{ingested} / kg _{released}].....	266
Table 9-7:	Values of the parameter sensitivity evaluation measure (term in brackets of Eq. (9-1)) for pan-European emissions of chromium to	

	air in 1990 for time horizons of 25 years and time-integrated (emissions only take place in the first year)	268
Table 9-8:	Contribution of the different food items to the Intake Fraction (last row) of chromium for ingestion exposures after 25 years and time-integrated according to the sensitivity cases with respect to pH variability (pan-European emissions to air in 1990, only taking place in the first year).....	272
Table 10-1:	Characteristics of the investigated power plants	278
Table 10-2:	Ranges of trace element concentrations in coals of different origin as quoted in Joint Research Centre of the European Commission (2003) [mg/kg]	279
Table 10-3:	Theoretical emission ranges of the investigated trace elements for the respective facilities [t/yr]	279
Table 10-4:	Effective Intake Fractions due to inhalation of selected trace elements for a one year pulse emission into air at different sites [kg _{inhaled} per kg _{released}].....	280
Table 10-5:	Disability Adjusted Life Years (DALYs) per kilogram of selected trace elements released due to cancer and non-cancer effects upon inhalation exposure caused by a one year pulse emission from selected power plant sites [years lost-equivalents per kg _{released}]	281
Table 10-6:	Damage factors due to inhalation for a one year pulse emission from different sites discounted at a rate of 0 and 3 % [€ ₂₀₀₀ per kg _{released}]	283
Table 10-7:	Ranges of quantifiable external costs discounted at 0 % due to inhalation of selected trace elements and in total caused by a one year pulse emission from different sites (variable units, base year 2000)	284
Table 10-8:	Ranges of quantifiable external costs discounted at 3 % due to inhalation of selected trace elements and in total caused by a one year pulse emission from different sites (variable units, base year 2000)	285
Table 10-9:	Time-integrated Disability Adjusted Life Years (DALYs) per kilogram of trace element released due to cancer and non-cancer effects upon ingestion exposure caused by a one year pulse emission from single sites [years lost-equivalents per kg _{released}].....	292
Table 10-10:	Damage factors due to ingestion for a one year pulse emission according to emissions from the Belgian power plant [€ ₂₀₀₀ per kg _{released}]	293
Table 10-11:	Damage factors due to ingestion for a one year pulse emission according to emissions from the French power plant [€ ₂₀₀₀ per kg _{released}]	294

Table 10-12: Damage factors due to ingestion for a one year pulse emission according to emissions from the German power plant [€_{2000} per $\text{kg}_{\text{released}}$]	295
Table 10-13: Damage factors due to ingestion for a one year pulse emission according to emissions from the power plant in the UK [€_{2000} per $\text{kg}_{\text{released}}$]	296
Table 10-14: Ranges of quantifiable external costs due to ingestion caused by a one year pulse emission of selected trace elements and in total according to theoretical minimum and maximum emission values from the Belgian power plant (variable units).....	297
Table 10-15: Ranges of quantifiable external costs due to ingestion caused by a one year pulse emission of selected trace elements and in total according to theoretical minimum and maximum emission values from the French power plant (variable units).....	298
Table 10-16: Ranges of quantifiable external costs due to ingestion caused by a one year pulse emission of selected trace elements and in total according to theoretical minimum and maximum emission values from the German power plant (variable units)	299
Table 10-17: Quantifiable external costs due to ingestion caused by a one year pulse emission of selected trace elements and in total according to theoretical minimum and maximum emission values from the power plant in the UK (variable units)	300
Table 11-1: Arsenic concentrations at steady-state in the Hron River catchment in central Slovakia according to the pan-European emission scenario for 1990 (continuous releases).....	306
Table 11-2: Effective Intake Fractions [$\text{kg}_{\text{inhaled}}$ per $\text{kg}_{\text{released}}$] and resulting cancer and non-cancer associated Disability Adjusted Life Years (DALYs) per kilogram of a trace element released [years lost-equivalents per $\text{kg}_{\text{released}}$] due to inhalation caused by a one year pulse emission into air according to the pan-European emission scenario for 1990	307
Table 11-3: Damage factors due to inhalation for a one year pulse emission according to the pan-European emission scenario for 1990 [€_{2000} per $\text{kg}_{\text{released}}$]	308
Table 11-4: Quantifiable external costs due to inhalation of selected trace elements and in total caused by a one year pulse emission according to the pan-European emission scenario for 1990 [$10^6 \text{€}_{2000}/\text{yr}$].....	309
Table 11-5: Time-integrated Disability Adjusted Life Years (DALYs) per kilogram of trace element released due to cancer and non-cancer effects upon ingestion exposure caused by a one year pulse emission ac-	

	cording to the pan-European emission scenario for 1990 [years lost-equivalents per kg _{released}]	312
Table 11-6:	Damage factors due to ingestion for a one year pulse emission according to the pan-European emission scenario for 1990 [€ ₂₀₀₀ per kg _{released}]	314
Table 11-7:	Quantifiable external costs due to ingestion of selected pollutants and in total caused by a one year pulse emission according to the pan-European emission scenario for 1990 [10 ⁶ € ₂₀₀₀ /yr]	315
Table 11-8:	Total trace element emissions in Europe in 1990 and 2000 estimated according to Droste-Franke et al. (2003) and ESPREME (2004), respectively	316
Table A-1:	Parameter needs for the assessment of particle deposition to above-ground produce that are neither related to substance nor to plant characteristics (like plant biomass, time until harvest).....	425
Table B-1:	Adjusted representation of catchments as given by the HYDRO1k dataset (EROS Data Center, 1996)	457
Table B-2:	Translation of CORINE land uses (European Environment Agency, 2000) into WATSON land uses	468
Table B-3:	Translation of USGS land uses (EROS Data Center et al., 2000) into WATSON land uses	470
Table B-4:	Deriving stream freshwater volumes per catchment area depending on the continent (drainage areas and discharges in the peripheral and central regions of the world taken from Baumgartner and Liebscher, 1990).....	475
Table B-5:	Organic carbon content and pH values for compartments other than permeable soils as used by WATSON	477
Table B-6:	Classes of pH values as given by Batjes (1996) and assigned representative single pH values	480
Table B-7:	Organic carbon classes as given by Batjes (1996) and assigned organic carbon reservoir values [kg _{carbon} /m ²]	480
Table B-8:	Reported erosion rates in multimedia models	488
Table B-9:	Characteristics of suspended matter as reported for some multimedia models and implications	491
Table B-10:	Characteristics of sediment solids as reported for some multimedia models and implications	492
Table B-11:	Characteristics of solids in the freshwater environment as used in the presented methodology	493
Table B-12:	Water discharge, area, sediment discharge and volume fractions of transported sediment for several rivers in the geographical scope of WATSON as compiled in Milliman and Syvitski (1992)	495

Table B-13: Relationships between the different process rates active in the particle mass balance for surface freshwater as assumed in this study	501
Table B-14: Reported invariant (organic) particle mass balances in surface freshwater bodies of non-site-specific multimedia models	504
Table B-15: Reported invariant (organic) particle mass balances in surface freshwater bodies of site-specific multimedia models	505
Table B-16: Annual national production of different produce [kt/yr]	512
Table B-17: Correction of country total receptor values according to area covered by WATSON	517
Table B-18: Difference between cow milk production and overall milk production and assumed share of cow milk production with respect to the overall milk production	520
Table B-19: Degree of self-supply in Europe with respect to the food groups considered in the exposure assessment; derived based on (Food and Agriculture Organization of the United Nations - Statistics Division, 2002a; Food and Agriculture Organization of the United Nations - Statistics Division, 2003).....	523
Table B-20: Ingestion of feed and soil particles by farm animals ($ING_{\text{feed or soil}}$) [kg DW/capita/s] according to United States - Environmental Protection Agency (1998)	524
Table B-21: Ingestion of forage by farm animals (ING_{forage}) [kg DW/capita/s] according to International Atomic Energy Agency (2001)	525
Table B-22: Mass fraction of food dry matter ($fr_{w_{\text{solid phase/bulk}}}$) [kg _{food} DW per kg _{food} FW]	526
Table B-23: Mass fraction of grains fed to farm animals consisting of wheat ($fr_{w_{\text{wheat/total grain}}}$) [-]	526
Table C-1: pH-dependent solid-water partitioning coefficients (K_d) [kg/kg _{solid phase} per kg/m ³ _{aqueous phase}].....	530
Table C-2: Mass fraction of a substance contained in food leading to an effect ($fr_{w_{\text{effective/total}}}$) [-]	534
Table C-3: Bioconcentration factors (BCF) aqueous phase-freshwater fish [m ³ /kg FW] (source: United States - Environmental Protection Agency, 1998)	535
Table C-4: Bioconcentration factors (BCF) for different vegetal produce [mol/kg plant DW per mol/kg soil DW] (source: United States - Environmental Protection Agency, 1998).....	536
Table C-5: Biotransfer factors (BTF) relating daily pollutant intake to contents in animal produce [s·capita/kg FW] (source: United States - Environmental Protection Agency, 1998).....	538

Table C-6:	Empirical correction factor ($\text{emp}_{\text{BCF, root crop}}$) for equilibrium uptake by belowground produce depending on the substance's octanol-water partitioning coefficient (K_{ow}) [-] (source: United States - Environmental Protection Agency, 1998, p. 5-35)	539
Table C-7:	Mass fraction adhering to aboveground exposed produce during wet deposition ($\text{fr}_{\text{w adhere/wet deposition}}$) [-] (source: United States - Environmental Protection Agency, 1998, Table B-2-7, p. B-78; value for cations and insoluble particles).....	539
Table C-8:	Parameter values used for the exposure assessment according to International Atomic Energy Agency (2001)	541
Table C-9:	Reported arsenic concentrations in environmental media and foodstuff.....	543
Table C-10:	Reported cadmium concentrations in environmental media and foodstuff.....	547
Table C-11:	Reported chromium concentrations in environmental media and foodstuff.....	551
Table C-12:	Reported lead concentrations in environmental media and foodstuff	554
Table D-1:	Symbols and corresponding unique units used.....	560
Table D-2:	Symbols used to show degrees of freedom. Symbols occur in parentheses or as indices	563
Table D-3:	Compartment acronyms employed.....	563