

## LIST OF FIGURES

<b>FIGURE 1.1</b> Pattern of rainfall acidity in Europe based on EMEP measurements (EMEP, 1988).....	2
<b>FIGURE 1.2</b> Intensity of defoliation for all species, or conifers only in 1988. Percentage of trees in classes 1-4 (>10% defoliation).....	4
<b>FIGURE 1.3</b> Cause and effect chain studied in the DPA (Heij and Schneider, 1991).....	5
<b>FIGURE 1.4</b> Schematic representation of deposition.....	6
<b>FIGURE 2.1</b> Emissions of sulphur dioxide, nitrogen oxides and ammonia in Europe from 1900 to 1990 (mton a <sup>-1</sup> ) compiled by De Leeuw (1994).....	27
<b>FIGURE 2.2</b> Emissions of sulphur dioxide, nitrogen oxides and ammonia in USA from 1900 to 1990 (mton a <sup>-1</sup> ) (NAPAP, 1992).....	27
<b>FIGURE 2.3</b> Annual average total SO <sub>2</sub> , NO <sub>x</sub> and NH <sub>3</sub> emissions in the Netherlands .....	29
<b>FIGURE 2.4</b> Atmospheric chemistry for trace gases in the troposphere.....	31
<b>FIGURE 2.5</b> Schematic representation of the main circulation types above western and central Europe. The location of the high pressure area is indicated by dots. The arrows give the tracking direction of depressions as well as the mean direction of the wind in the higher air layers. Generally, the characters represent the main direction of the circulation (e.g. SE = southeast); Ww = bending west circulation, BM = high pressure area above Atlantic ocean - central Europe, HM = high pressure area above central Europe, TM = low pressure area above central Europe (after Van der Ham, 1977). .....	37
<b>FIGURE 2.6</b> Daily variation of the boundary-layer development (Stull, 1988).....	38
<b>FIGURE 2.7</b> Distribution of annual median SO <sub>2</sub> (A) and NO <sub>2</sub> (B) concentrations in the Netherlands in 1993 (µg m <sup>-3</sup> ).....	42
<b>FIGURE 2.8</b> Distribution of annual average NH <sub>3</sub> concentrations in the Netherlands in 1993 (µg m <sup>-3</sup> ).....	43
<b>FIGURE 2.9</b> Annual country average of 50-percentile SO <sub>2</sub> , NO <sub>2</sub> , NO <sub>x</sub> and O <sub>3</sub> concentrations, and annual average NH <sub>3</sub> (µg m <sup>-3</sup> ) in 1977 - 1993 based on LML measurements (RIVM, 1994) and TREND model calculations (Van Jaarsveld, 1995).....	44

<b>FIGURE 2.10</b> Annual mean concentration of SO <sub>2</sub> (above) and SO <sub>4</sub> <sup>2-</sup> aerosol (below) in 1993 in Europe as calculated by Tuovinen <i>et al.</i> (1994) using the EMEP-LTRAP model .....	45
<b>FIGURE 2.10</b> ( <i>continued</i> ) Annual mean concentration of NO <sub>x</sub> (above) and HNO <sub>3</sub> (below) in 1993 in Europe as calculated by Tuovinen <i>et al.</i> (1994) using the EMEP-LTRAP model .....	46
<b>FIGURE 2.10</b> ( <i>continued</i> ) Annual mean concentration of NO <sub>3</sub> <sup>-</sup> aerosol (above) and NH <sub>3</sub> (below) in 1993 in Europe as calculated by Tuovinen <i>et al.</i> (1994) using the EMEP-LTRAP model.....	47
<b>FIGURE 2.10</b> ( <i>continued</i> ) Annual mean concentration of NH <sub>4</sub> <sup>+</sup> aerosol in 1993 in Europe as calculated by Tuovinen <i>et al.</i> (1994) using the EMEP-LTRAP model .....	48
<b>FIGURE 3.1</b> Wet deposition processes. ....	50
<b>FIGURE 3.2</b> The dry deposition process.....	55
<b>FIGURE 3.3</b> Schematic representation of four transport mechanisms for particles through the quasi-laminar boundary layer (after Davidson and Wu, 1990). Shaded areas indicate the laminar boundary layer. ....	61
<b>FIGURE 3.4</b> Relationship between the deposition velocity of particles and their mass median diameter for an Eucalyptus forest canopy (after Slinn, 1982). The dashed line indicates the sedimentation velocity (i.e. where the deposition rate is a linear function of the fall velocity of the particles). Canopy height = 27.4 m; friction velocity = 75 cm s <sup>-1</sup> ; roughness length = 1.86 m; zero plain displacement height = 21.6 m; particle density = 1 g cm <sup>-3</sup> . ....	62
<b>FIGURE 3.5</b> Illustration of instrumentation used for flux gradient measurements.....	70
<b>FIGURE 3.5</b> ( <i>continued</i> ) Illustration of instrumentation used for flux gradient measurements.....	71
<b>FIGURE 5.1</b> Frequency distribution of deposition on a small scale (5 x 5 km) within an EMEP grid (150 x 150 km) for nitrogen compounds. Using EMEP model results together with the appropriate filtering factor, we estimate total nitrogen deposition to this grid at 1650 mol ha <sup>-1</sup> a <sup>-1</sup> . ....	115
<b>FIGURE 5.2</b> Distribution of the acidification areas over the Netherlands. ....	117
<b>FIGURE 5.3</b> The spatial distribution of the dry deposition of SO <sub>4</sub> <sup>2-</sup> , NO <sub>3</sub> <sup>-</sup> , NH <sub>4</sub> <sup>+</sup> and total potential acid in 1993 (mol ha <sup>-1</sup> a <sup>-1</sup> ).....	121
<b>FIGURE 5.4</b> Values for z <sub>0</sub> as a function of wind direction sector (30 <sup>0</sup> ) and season for the Steeg station (103). ....	124
<b>FIGURE 5.5</b> Values for z <sub>0</sub> as a function of wind direction sector (30 <sup>0</sup> ) for the Cabauw station (620) estimated by Beljaars (1988) and in this study for summer (A) and winter (B) months.....	125
<b>FIGURE 5.6</b> 1 x 1 km grid average z <sub>0</sub> values in the Netherlands: A: 0-60 cm, B: >50 cm. .	127
<b>FIGURE 5.7</b> The spatial distribution of the dry deposition of SO <sub>x</sub> , NO <sub>y</sub> , NH <sub>x</sub> and total potential acid in 1993 (mol ha <sup>-1</sup> a <sup>-1</sup> ).....	129
<b>FIGURE 5.8</b> Distribution of dry deposition SO <sub>x</sub> to forest, grassland and water in 1993 in the Netherlands. ....	130

---

<b>FIGURE 5.8</b> ( <i>continued</i> ) Distribution of dry deposition of NO <sub>y</sub> and NH <sub>x</sub> to forest, grassland and water in 1993 in the Netherlands. ....	131
<b>FIGURE 5.9</b> Maps of the dry deposition of Na <sup>+</sup> (A), K <sup>+</sup> (B), Mg <sup>2+</sup> (C) and Ca <sup>2+</sup> (D) in the Netherlands (mol ha <sup>-1</sup> a <sup>-1</sup> ). ....	133
<b>FIGURE 5.10</b> Dry deposition distribution of Na <sup>+</sup> , K <sup>+</sup> , Mg <sup>2+</sup> and Ca <sup>2+</sup> for forest, grassland and water. ....	134
<b>FIGURE 5.10</b> ( <i>continued</i> ) Distribution of dry deposition Mg <sup>2+</sup> and Ca <sup>2+</sup> to forest, grassland and water. ....	135
<b>FIGURE 5.11</b> Variation in total SO <sub>x</sub> , NO <sub>y</sub> , NH <sub>x</sub> and potential acid deposition between 1980 and 1993 (mol ha <sup>-1</sup> a <sup>-1</sup> ). ....	136
<b>FIGURE 5.12</b> The spatial distribution of the total deposition of SO <sub>x</sub> , NO <sub>y</sub> , NH <sub>x</sub> and total potential acid in 1980 (mol ha <sup>-1</sup> a <sup>-1</sup> ). ....	137
<b>FIGURE 5.13</b> The spatial distribution of the total deposition of SO <sub>x</sub> , NO <sub>y</sub> , NH <sub>x</sub> and total potential acid in 1993 (mol ha <sup>-1</sup> a <sup>-1</sup> ). ....	138
<b>FIGURE 5.14</b> Contribution of different countries to the deposition in the Netherlands (A) 1980 and (B) 1993. ....	142
<b>FIGURE 5.15</b> Source category contribution to the deposition in the Netherlands (A) 1980 and (B) 1990. ....	143
<b>FIGURE 5.16</b> The spatial distribution of the deposition of Na, K, Mg <sup>2+</sup> and Ca in 1993 (mol ha <sup>-1</sup> a <sup>-1</sup> ). ....	145
<b>FIGURE 5.17</b> Outline of method to estimate local scale deposition fluxes. ....	147
<b>FIGURE 5.18</b> Location of wet deposition monitoring sites in Europe. ....	148
<b>FIGURE 5.19</b> Wet deposition of SO <sub>4</sub> <sup>2-</sup> in Europe on a 50 x 50 km basis in 1989 in mol ha <sup>-1</sup> a <sup>-1</sup> (Van Leeuwen <i>et al.</i> , 1995). ....	151
<b>FIGURE 5.20</b> Wet deposition of NO <sub>3</sub> <sup>-</sup> in Europe on a 50 x 50 km basis in 1989 in mol ha <sup>-1</sup> a <sup>-1</sup> (Van Leeuwen <i>et al.</i> , 1995). ....	151
<b>FIGURE 5.21</b> Wet deposition of NH <sub>4</sub> <sup>+</sup> in Europe on a 50 x 50 km basis in 1989 in mol ha <sup>-1</sup> a <sup>-1</sup> (Van Leeuwen <i>et al.</i> , 1995). ....	152
<b>FIGURE 5.22</b> Wet deposition of H <sup>+</sup> in Europe on a 50 x 50 km basis in 1989 in mol ha <sup>-1</sup> a <sup>-1</sup> (Van Leeuwen <i>et al.</i> , 1995). ....	152
<b>FIGURE 5.23</b> Wet deposition of total base cations (Ca <sup>2+</sup> , Mg <sup>2+</sup> and K <sup>+</sup> ) in Europe on a 50 x 50 km basis in 1989 in mol ha <sup>-1</sup> a <sup>-1</sup> (Van Leeuwen <i>et al.</i> , 1995). ....	153
<b>FIGURE 5.24</b> Concentration of Na <sup>+</sup> in Europe on a 50 x 50 km basis in 1989 in μmol l <sup>-1</sup> (Van Leeuwen <i>et al.</i> , 1995). ....	154
<b>FIGURE 5.25</b> Concentration of Cl <sup>-</sup> in Europe on a 50 x 50 km basis in 1989 in μmol l <sup>-1</sup> (Van Leeuwen <i>et al.</i> , 1995). ....	155
<b>FIGURE 5.26</b> Concentration of Mg <sup>2+</sup> in Europe on a 50 x 50 km basis in 1989 in μmol l <sup>-1</sup> (Van Leeuwen <i>et al.</i> , 1995). ....	155
<b>FIGURE 5.27</b> Concentration of Ca <sup>2+</sup> in Europe on a 50 x 50 km basis in 1989 in μmol l <sup>-1</sup> (Van Leeuwen <i>et al.</i> , 1995). ....	156

---

---

<b>FIGURE 5.28</b> Concentration of $K^+$ in Europe on a 50 x 50 km basis in 1989 in $\mu\text{mol l}^{-1}$ (Van Leeuwen <i>et al.</i> , 1995).....	156
<b>FIGURE 5.29</b> Land-use map on a $1/6^0 \times 1/6^0$ grid of Europe (Van Velde, 1994).....	157
<b>FIGURE 5.30</b> Roughness length map (cm) for the summer season (A) and winter season (B) on a $1/6^0 \times 1/6^0$ grid of Europe (Van Pul <i>et al.</i> , 1995). .....	158
<b>FIGURE 5.31</b> Dry deposition of $\text{SO}_x$ in Europe on a $1/6^0 \times 1/6^0$ scale in $\text{mol ha}^{-1} \text{a}^{-1}$ (Van Pul <i>et al.</i> , 1995).....	159
<b>FIGURE 5.32</b> Dry deposition of $\text{NO}_y$ in Europe on a $1/6^0 \times 1/6^0$ scale in $\text{mol ha}^{-1} \text{a}^{-1}$ (Van Pul <i>et al.</i> , 1995).....	160
<b>FIGURE 5.33</b> Dry deposition of $\text{NH}_x$ in Europe on a $1/6^0 \times 1/6^0$ scale in $\text{mol ha}^{-1} \text{a}^{-1}$ (Van Pul <i>et al.</i> , 1995).....	160
<b>FIGURE 5.34</b> Dry deposition of $\text{Na}^+$ in Europe on a $1/6 \times 1/6^0$ scale in $\text{mol ha}^{-1} \text{a}^{-1}$ .....	161
<b>FIGURE 5.35</b> Dry deposition of $\text{Ca}^{2+}$ in Europe on a $1/6 \times 1/6^0$ scale in $\text{mol ha}^{-1} \text{a}^{-1}$ .....	162
<b>FIGURE 5.36</b> Dry deposition of $\text{Mg}^{2+}$ in Europe on a $1/6 \times 1/6^0$ scale in $\text{mol ha}^{-1} \text{a}^{-1}$ .....	162
<b>FIGURE 5.37</b> Dry deposition of $K^+$ in Europe on a $1/6 \times 1/6^0$ scale in $\text{mol ha}^{-1} \text{a}^{-1}$ .....	163
<b>FIGURE 5.38</b> Dry deposition of $K^+$ in Europe on a $1/6 \times 1/6^0$ scale in $\text{mol ha}^{-1} \text{a}^{-1}$ .....	163
<b>FIGURE 5.39</b> Total deposition of $\text{SO}_x$ in Europe on a $1/6 \times 1/6^0$ scale in $\text{mol ha}^{-1} \text{a}^{-1}$ (Van Pul <i>et al.</i> , 1995).....	164
<b>FIGURE 5.40</b> Total deposition of $\text{NO}_y$ in Europe on a $1/6 \times 1/6^0$ scale in $\text{mol ha}^{-1} \text{a}^{-1}$ (Van Pul <i>et al.</i> , 1995).....	165
<b>FIGURE 5.41</b> Total deposition of $\text{NH}_x$ in Europe on a $1/6 \times 1/6^0$ scale in $\text{mol ha}^{-1} \text{a}^{-1}$ (Van Pul <i>et al.</i> , 1995).....	165
<b>FIGURE 5.42</b> Total deposition of potential acid in Europe on a $1/6 \times 1/6^0$ scale in $\text{mol ha}^{-1} \text{a}^{-1}$ (Van Pul <i>et al.</i> , 1995).....	166
<b>FIGURE 5.43</b> Emission of $\text{SO}_x$ , $\text{NO}_y$ and $\text{NH}_x$ per country or region in Europe in 1993 expressed in $\text{mol H}^+ \text{ha}^{-1} \text{a}^{-1}$ (Tuovinen <i>et al.</i> , 1994).....	168
<b>FIGURE 5.44</b> Deposition of $\text{SO}_x$ , $\text{NO}_y$ and $\text{NH}_x$ per country or region in Europe in 1993 expressed in $\text{mol H}^+ \text{ha}^{-1} \text{a}^{-1}$ (Tuovinen <i>et al.</i> , 1994).....	169
<b>FIGURE 5.45</b> Emission and deposition of potential acid per country or region in Europe in 1993 in $\text{mol H}^+ \text{ha}^{-1} \text{a}^{-1}$ (Tuovinen <i>et al.</i> , 1994).....	170
<b>FIGURE 5.46</b> Contribution (%) of different source categories to the total potential acid deposition for different countries in Europe.....	171
<b>FIGURE 5.47</b> Total deposition of $\text{Na}^+$ in Europe on a $1/6 \times 1/6^0$ scale in $\text{mol ha}^{-1} \text{a}^{-1}$ .....	172
<b>FIGURE 5.48</b> Total deposition of $\text{Ca}^{2+}$ in Europe on a $1/6 \times 1/6^0$ scale in $\text{mol ha}^{-1} \text{a}^{-1}$ .....	173
<b>FIGURE 5.49</b> Total deposition of $\text{Mg}^{2+}$ in Europe on a $1/6 \times 1/6^0$ scale in $\text{mol ha}^{-1} \text{a}^{-1}$ .....	173
<b>FIGURE 5.50</b> Total deposition of $K^+$ in Europe on a $1/6 \times 1/6^0$ scale in $\text{mol ha}^{-1} \text{a}^{-1}$ .....	174
<b>FIGURE 5.51</b> Total deposition of base cations ( $\text{Ca}^{2+} + \text{Mg}^{2+} + \text{K}^+$ ) in Europe on a $1/6 \times 1/6^0$ scale in $\text{mol ha}^{-1} \text{a}^{-1}$ .....	174
<b>FIGURE 5.52</b> Wet deposition of ammonium at Rothamsted (UK) and Montsouris (France) (in $\text{kg N ha}^{-1} \text{a}^{-1}$ ).....	178

---

**FIGURE 5.52** (*continued*) Wet deposition of nitrate at Rothamsted (UK) and Montsouris (France) (in  $\text{kg N ha}^{-1} \text{ a}^{-1}$ )..... 179

**FIGURE 5.53** Precipitation concentrations of sulphur measured over several years ( $\text{mg SO}_3 \text{ l}^{-1}$ )..... 181

**FIGURE 5.54** Changes in  $\text{Na}^+$ ,  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  concentrations in precipitation ( $\text{mg l}^{-1}$ ) measured at Leiduin near the Dutch coast. .... 183

**FIGURE 5.55** Horizontal gradient of wet deposition from the coast land inward as reported by Leeftang (1938). Data are averaged over 5 years (1932 - 1937)..... 184

**FIGURE 5.56** Time series of  $\text{SO}_4^{2-}$  concentration measurements in precipitation made at several sites in the Netherlands ( $\text{mg l}^{-1}$ )..... 185

**FIGURE 5.56** (*continued*) Time series of  $\text{NO}_3^-$  concentration measurements in precipitation made at several sites in the Netherlands ( $\text{mg l}^{-1}$ )..... 186

**FIGURE 5.57** Time series of wet deposition measurements made at Witteveen: (A)  $\text{SO}_4^{2-}$ , (B)  $\text{NO}_3^-$  and (C)  $\text{NH}_4^+$  ( $\text{mol ha}^{-1} \text{ a}^{-1}$ ). .... 187

**FIGURE 5.57** Time series of wet deposition measurements made at Witteveen: (A)  $\text{SO}_4^{2-}$ , (B)  $\text{NO}_3^-$  and (C)  $\text{NH}_4^+$  ( $\text{mol ha}^{-1} \text{ a}^{-1}$ ). .... 188

**FIGURE 5.58** Relative change between 1955 and 1990 in measured concentrations of sulphur in air, aerosol and precipitation, and in estimated sulphur emissions in the Netherlands and in Europe, and in ammonia emissions in the Netherlands. .... 190

**FIGURE 5.59** Temporal variation in wet and dry deposition of sulphur in the Netherlands ( $\text{mol ha}^{-1} \text{ a}^{-1}$ ). .... 191

**FIGURE 5.60** Ratio of  $\text{NH}_4^+$  over ( $\text{SO}_4^{2-}$  plus  $\text{NO}_3^-$ ) in precipitation in the Netherlands..... 192

**FIGURE 6.1** Comparison of modelled  $V_d$  ( $\text{m s}^{-1}$ ) with  $V_d$  obtained from measurements: A. Deciduous forest; B. Coniferous forest; C. Grassland and D. Heathland. Solid dots represent average modelled values for class-average measured values. The line represents the 1:1 ratio. Small overbars represent measured class averages  $\pm$  SD, while large overbars represent modelled averages  $\pm$  SD. .... 199

**FIGURE 6.2.** Examples of time series of modelled and measured  $V_d$ : A. Deciduous forest; B. Coniferous forest..... 203

**FIGURE 6.2** (*continued*). Examples of time series of modelled and measured  $V_d$ : C. Grassland and D. Heathland. .... 204

**FIGURE 6.3** Histograms of modelled (open columns) and measured (dark columns)  $V_d$  ( $\text{m s}^{-1}$ ): A. Deciduous forest; B. Coniferous forest. .... 206

**FIGURE 6.3** (*continued*) Histograms of modelled (open columns) and measured (dark columns)  $V_d$  ( $\text{m s}^{-1}$ ): C. Grassland and D. Heathland ..... 207

**FIGURE 6.4** Throughfall estimates compared to DEADM results for  $\text{SO}_x$  (A),  $\text{NO}_y$  (B),  $\text{NH}_x$  (C) and total potential acid (D) ( $\text{mol ha}^{-1} \text{ a}^{-1}$ ) for 21 sites. The 1:1 line is also shown. .... 221

**FIGURE 6.5** Throughfall estimates compared to DEADM results for  $\text{SO}_x$  (A),  $\text{NO}_y$  (B),  $\text{NH}_x$  (C) and total potential acid (D) ( $\text{mol ha}^{-1} \text{ a}^{-1}$ ) for 30 sites. The 1:1 line is also shown. .... 222

---

**FIGURE 6.6** Net throughfall estimates compared to DEADM dry deposition estimates for 30 sites for Na<sup>+</sup> (A), Mg<sup>2+</sup> (B), Ca<sup>2+</sup> (C) and K<sup>+</sup> (D) (mol ha<sup>-1</sup> a<sup>-1</sup>). The 1:1 line is also shown. .... 223

**FIGURE 6.7** Comparison of measured concentrations in Speulder forest (30 m) and those obtained from the model's calculations (50 m). .... 227

**FIGURE 6.8** Comparison of monthly  $V_d$  values calculated with DEADM (50-m height) and measured in Speulder forest (36-m height) (m s<sup>-1</sup>). .... 227

**FIGURE 6.9** Comparison of TREND model results with estimates from this study (mol ha<sup>-1</sup> a<sup>-1</sup>). The 1:1 line is shown for comparison. .... 229

**FIGURE 6.10** Comparison of the country average calculated dry deposition of sulphur using EMEP and EDACS (mol ha<sup>-1</sup> a<sup>-1</sup>). .... 245

**FIGURE 6.11** Distribution of the numbered (pollution) regions in Europe in Table 6.13. .... 246

**FIGURE 7.1** Time series of parameterized and measured  $R_c$  for SO<sub>2</sub> at Elspeetsche Veld. .... 258

**FIGURE 7.2** Averaged  $R_c$  for NO<sub>2</sub> over heathland (Elspeetsche Veld) in autumn 1989 (Duyzer *et al.*, 1991). .... 260

**FIGURE 7.3** Annual average diurnal variations of  $F$  and  $V_d$  for NH<sub>3</sub>. .... 261

**FIGURE 7.4** Surface resistance for SO<sub>2</sub> and NH<sub>3</sub>, relative humidity and surface wetness indicator for the Elspeetsche Veld site, 14-15 September 1991 (Erisman and Wyers, 1993). .... 262

**FIGURE 7.5** Location of the study area with the 30 throughfall measurement sites and five bulk precipitation measurement sites used to study the impact of canopy structure on dry deposition amounts. .... 270

**FIGURE 7.6** Location of the study area with the eight forest edge measurement sites and four bulk precipitation measurement sites used to study the impact of edge effects on dry deposition amounts. .... 271

**FIGURE 7.7** An example of a measurement site showing the area (10 x 10 m) around the throughfall gutter in which tree and stand structure characteristics were determined. The throughfall gutter, indicated as a black bar, is located in the centre of the plot. Grey areas mark the vertical crown projection of individual trees, whereas stem bases are indicated as black circles. The thick line indicates the boundary of the 100 m<sup>2</sup> plot. .... 275

**FIGURE 7.8** Relationship between winter net throughfall fluxes of SO<sub>4</sub><sup>2-</sup> and the aerodynamic roughness length of the canopy. Douglas fir is indicated by squares, Scots pine by plus signs and oak by dots. The following relationship is valid:  $[NTF-SO_4^{2-}] = 344.9*[z_0] + 201.3$  ( $R=0.84$ ;  $p<0.001$ ). .... 286

**FIGURE 7.9** Relationship between winter net throughfall fluxes of NO<sub>3</sub><sup>-</sup> and the aerodynamic roughness length of the canopy. Douglas fir is indicated by squares, Scots pine by plus signs and oak by dots. The following relationship is valid:  $[NTF-NO_3^-] = 59.6*[z_0] + 21.5$  ( $R=0.84$ ;  $p<0.001$ ). .... 286

**FIGURE 7.10** Relationship between winter net throughfall fluxes of NH<sub>4</sub><sup>+</sup> and the aerodynamic roughness length of the canopy. Douglas fir is indicated by squares, Scots

pine by plus signs and oak by dots. The following relationship is valid:  $[NTF-NH_4^+]=431.0*[z_o] +65.5$  ( $R=0.88$ ;  $p<0.001$ )..... 287

**FIGURE 7.11** Winter net throughfall fluxes of  $SO_4^{2-}$  predicted from linear regression models with the roughness length of the canopy, compared to measured winter net throughfall fluxes. Douglas fir is indicated by squares, Scots pine by plus signs and oak by dots... 289

**FIGURE 7.12** Winter net throughfall fluxes of  $NO_3^-$  predicted from linear regression models with the roughness length of the canopy, compared to measured winter net throughfall fluxes. Douglas fir is indicated by squares, Scots pine by plus signs and oak by dots... 289

**FIGURE 7.13** Winter net throughfall fluxes of  $NH_4^+$  predicted from linear regression models with the roughness length of the canopy, compared to measured winter net throughfall fluxes. Douglas fir is indicated by squares, Scots pine by plus signs and oak by dots... 290

**FIGURE 7.14** Net throughfall flux gradients of  $SO_4^{2-}$ ,  $NO_3^-$  and  $NH_4^+$  for the eight forest edges. Distance to edge divided by edge height is  $x/h$ ..... 293

**FIGURE 7.14 (continued)** Net throughfall flux gradients of  $Na^+$ ,  $Cl^-$  and  $Mg^{2+}$  for the eight forest edges. Distance to edge divided by edge height is  $x/h$ ..... 294

**FIGURE 7.14 (continued)** Net throughfall flux gradients of  $Ca^{2+}$  and  $K^+$ , and gradients of the ratio between throughfall volume and precipitation volume for the eight forest edges. Distance to edge divided by edge height is  $x/h$ ..... 295

**FIGURE 7.15** ‘Ideal’ net throughfall flux gradient in a forest edge..... 296

**FIGURE 7.16** The percentage to which a particular wind direction contributed to the total pollutant dose of  $SO_2$ ,  $SO_4^{2-}$ ,  $NO_2$ ,  $NO$ ,  $NO_3^-$ , and  $NH_4^+$  during the measurements, respectively ..... 301

**FIGURE 7.16 (continued)** The percentage to which a particular wind direction contributed to the total pollutant dose of  $Na^+$ ,  $Cl^-$ ,  $Mg^{2+}$ ,  $Ca^{2+}$  and  $K^+$ , and total air supply during the measurements, respectively ..... 302

**FIGURE 7.17** The percentage to which different size classes contribute to the total number (and area) of forest complexes (left), and to the total number (and area) of individual forest stands in the Netherlands (right)..... 305

**FIGURE 7.18** Frequency distribution of forest stand heights in the Netherlands. The relatively high percentage observed for height class 0-2 m is the result of the relatively high fraction of timber felling areas and non-afforested silvicultural areas in the Netherlands..... 306

**FIGURE 7.19** Relationship between the part of the total area of Dutch forests influenced by edge effects and edge width. The thin line represents this relationship when only edges between forested areas and non-forested areas are taken into account..... 307

**FIGURE 7.20** Deposition research at the Speulder forest site. .... 315

**FIGURE 7.21** Eddy correlation  $u^*$  measurements at 30 m compared to those measured at 36 m ..... 317

**FIGURE 7.22** Eddy correlation  $H$  measurements at 30 m compared to those measured at 36 m. .... 318

**FIGURE 7.23** Daily variations of  $SO_2$  in  $R_c$  for dry and wet surface conditions..... 320

---

**FIGURE 7.24**  $R_c$  values for  $\text{SO}_2$  and  $\text{NH}_3$  from continuous measurements of vertical gradients over the Speulder forest ('Weather' provides information on precipitation: 0=dry; 10=rain). ..... 321

**FIGURE 7.25** Diurnal variation of  $V_d \text{NO}_2$  and  $V_d$  for stomatal uptake for selected measuring periods..... 323

**FIGURE 7.26** Parameterised values of  $V_d$  compared to  $V_d$  derived from  $\text{NH}_3$  gradients for selected measuring periods ( $R^2=0.25$  for positive values) ..... 324

**FIGURE 7.27** Experimentally determined  $V_d$  versus  $u^*$ .  $V_d$  values were estimated at the following heights: fog at 28 m;  $\text{NO}_3^-$  and  $\text{SO}_4^{2-}$  (TNO) at 35 m;  $\text{SO}_4^{2-}$  (ECN) at 22 m and  $^{214}\text{Pb}$  at 19 m. .... 330

**FIGURE 7.28** Summary of modelled deposition velocities versus  $u^*$ .  $V_d$  values were estimated at the following heights: fog at 28 m;  $\text{NO}_3^-$  and  $\text{SO}_4^{2-}$  (TNO) at 35 m;  $\text{SO}_4^{2-}$  (ECN) at 22 m and  $^{214}\text{Pb}$  at 19 m..... 336